

A POPULATION OF RADIO-LOUD NARROW LINE SEYFERT 1 GALAXIES WITH BLAZAR-LIKE PROPERTIES?

W. YUAN¹, H. Y. ZHOU^{2,3,4}, S. KOMOSSA⁴, X. B. DONG^{2,3}, T. G. WANG^{2,3}, H. L. LU^{2,3}, J. M. BAI¹

accepted for publication in ApJ

ABSTRACT

Blazars with strong emission lines were found to be associated mostly with broad-line type 1 Active Galactic Nuclei (AGN). Hitherto, evidence for blazars identified with Narrow Line Seyfert 1 (NLS1) AGN was limited to very few individual cases. Here we present a comprehensive study of a sample of 23 genuine radio-loud NLS1 galaxies which have the radio-loudness parameters, the ratio of radio (21 cm) to optical (4400Å) luminosity, greater than 100. The sample, drawn from the SDSS and FIRST, is homogeneous and the largest of this kind. A significant fraction of the sample objects show interesting radio to X-ray properties that are unusual to most of the previously known radio-loud NLS1 AGN, but are reminiscent of blazars. These include flat radio spectra, large amplitude flux and spectral variability, compact VLBI cores, very high brightness temperatures (10^{11-14} K) derived from variability, enhanced optical emission in excess of the normal ionising continuum, flat X-ray spectra, and blazar-like SEDs. We interpret them as evidence for the postulated blazar nature of these very radio-loud NLS1 AGN, which might possess at least moderately relativistic jets. We suggest that those steep spectrum radio-loud NLS1 AGN in the sample are of the same population but with their radio jets aligned at large angles to the lines-of-sight. Intrinsically, some of the objects have relatively low radio power and would have been classified as radio-intermediate AGN.

The black hole masses, estimated from the broad Balmer line width and luminosity, are within $10^{6-8} M_{\odot}$, and the inferred Eddington ratios are around unity. Unless the black hole masses are largely under-estimated, our result stretches the low mass end of the black holes of luminous, fast accreting radio-loud AGN to a smaller mass regime (the order of $10^6 M_{\odot}$) in the black hole mass–radio-loudness space where other normal AGN are seldom found. The results imply that radio-loud AGN may be powered by black holes with moderate masses ($\sim 10^{6-7} M_{\odot}$) accreting at high rates (Eddington ratios up to unity or higher). The host galaxies of a few nearby objects appear to be disk-like or merger; and some of the objects show imprints of young stellar populations in their SDSS spectra. We find that some of the objects, despite having strong emission lines, resemble high-energy peaked BL Lacs in their SED with the synchrotron component peaked at around the UV; such objects constitute an intriguingly high fraction of the sample. The radio sources of the sample are ubiquitously compact. They are smaller than at most several tens of kilo-parsecs, suggesting a possible link with compact steep-spectrum radio sources. Given the peculiarities of blazar-like NLS1 galaxies, questions arise as to whether they are plain downsizing extensions of normal radio-loud AGN, or whether they form a previously unrecognised population.

Subject headings: galaxies: active — galaxies: Seyfert — galaxies: jets — quasars: general — X-rays: galaxies — radio continuum: galaxies

1. INTRODUCTION

As a minority of AGN, radio-loud (RL) AGN differ from their radio-quiet (RQ) counterparts mainly in possessing prominent radio jets and/or lobes that produce strong radio radiation. In observations, a widely used division between RL and RQ AGN is the radio-loudness parameter $R \approx 10$, defined as the ratio between the radio 5 GHz to optical *B*-band luminosity (Kellermann et al. 1989; Stocke et al. 1992), though it is still a controversy whether R has a bimodal distribution below and above this value. RL AGN are an important laboratory to study the formation of relativistic jets, which is not yet understood so far (e.g. Blandford 2000; Celott & Blandford 2001; Meier 2003). Nevertheless, it is widely thought that radio-loudness (formation of radio jets) is possibly depending on the accretion rate/state (e.g. Ho 2002; Maccarone et

al. 2003; Greene et al. 2006; Körding et al. 2006), the spin of the black hole (e.g. Blandford 2000; Maraschi & Tavecchio 2003; Sikora et al. 2007), black hole mass (M_{BH} , e.g. Laor 2000; Lacy 2001; Dunlop et al. 2003; Mclure & Jarvis 2004; Metcalf & Magliocchetti 2006, but see Woo & Urry 2002 for a different view), and host galaxy morphology (e.g. Balmaverde & Capetti 2006; Capetti & Balmaverde 2006; Sikora et al. 2007), or a combination of some of these factors (e.g. Ballantyne 2007).

As an important sub-class of RL AGN, blazar is a collective term for BL Lac objects and flat-spectrum radio quasars (FSRQs). Blazars are characterised by flat radio spectra at above ~ 1 GHz, fast variability, high and variable polarization, superluminal motion, and high brightness temperatures (see e.g. Urry & Padovani 1995; Celott 2002, for reviews). They are now believed to be RL AGN with the orientation of relativistic jets close to the line-of-sight, and hence their non-thermal jet emission is highly Doppler boosted (Blandford & Rees 1978; Urry & Padovani 1995). Blazars have distinctive spectral energy distributions (SED) that are characterised by two broad humps in the $\log \nu - \log \nu f_{\nu}$ representation. The low-energy hump is commonly interpreted as synchrotron emission, while the high-energy one as inverse

Electronic address: wmy@ynao.ac.cn

¹ National Astronomical Observatories/Yunnan Observatory, Chinese Academy of Sciences, Kunming, Yunnan, P.O. BOX 110, P.R.China

² Center for Astrophysics, University of Science and Technology of China, Hefei, Anhui, 230026, P.R.China

³ Joint Institute of Galaxies and Cosmology, SHAO and USTC

⁴ Max-Planck-Institut für extraterrestrische Physik, Postfach 1312, 85741 Garching, Germany

Compton emission scattering off the same electron population which produces the synchrotron emission. Depending on the peak frequency of the synchrotron hump, BL Lac objects are divided into Low-energy-peaked BL Lacs (LBL) with the peak around IR–optical wavelengths and High-energy-peaked BL Lacs (HBL) with the peak around UV/soft X-ray energies (Padovani & Giommi 1995). Classical FSRQs have generally the peak frequencies of the synchrotron hump similar to or even lower than LBL. Interestingly, it was suggested that the whole blazar family can be arranged in a sequence from HBL to LBL and to FSRQ in the order of decreasing synchrotron peak frequencies and increasing source power—the so called blazar sequence⁵ (Fossati et al. 1998). However, exceptions were also reported in recent years as the presence of powerful FSRQs with HBL-like SEDs (though this remains a matter of controversy), whose synchrotron hump is peaked around the UV/soft X-ray band (see e.g. Padovani et al. 2003, and references therein). Following Perlman et al. (1998), we term FSRQs with HBL-like SED HFSRQs (High-energy-peaked FSRQs), and those with LBL-like SED—the classical FSRQs—LFSRQs (Low-energy-peaked FSRQs), as parallels to HBLs and LBLs.

Until recently, RL AGN (blazars) with strong broad emission lines were found to ubiquitously have the Balmer line widths greater than 2000 km s^{-1} in full width at half maximum (FWHM). It was found that the line width is correlated with the orientation of the radio axis, with narrower lines as well as stronger optical Fe II emission in objects with smaller viewing angles to the radio axis (e.g. Wills & Browne 1986; Jackson & Browne 1991). This was interpreted as arising from a disk-like emission line region. A marked absence of radio-loud quasars (RLQs) was noted below the 2000 km s^{-1} line-width cutoff (Wills & Browne 1986), that is the characteristic line-width range for narrow line Seyfert 1 galaxies—a sub-class of type 1 AGN in contrast to the classical Broad Line AGN (BLAGN). The situation has changed since the detection of radio emission from NLS1 galaxies.

Apart from narrow line-widths of the broad Balmer lines (originally defined as $< 2000 \text{ km s}^{-1}$, Osterbrock & Pogge 1985; Goodrich 1989), NLS1 galaxies also show other extreme properties compared to normal broad line AGN (see Komossa 2008, for a recent review), such as strong permitted optical/UV Fe II emission lines (Boroson & Green 1992; Grupe et al. 1999; Véron-Cetty et al. 2001), steep soft X-ray spectra (Wang et al. 1996; Boller et al. 1996; Grupe et al. 1998), and rapid X-ray variability (Leighly 1999; Komossa & Meerschweinchen 2000). Observational evidence suggests that NLS1 galaxies tend to have small black hole masses and high Eddington ratios (defined as the bolometric to the Eddington luminosity ratio), $R_{\text{Edd}} \equiv L_{\text{bol}}/L_{\text{Edd}} \approx 1$ (e.g. Boroson 2002; Collin & Kawaguchi 2004). In fact, they were found to locate at one extreme end, opposite to BLAGN, of eigenvector 1 of the correlation matrix which is believed to be driven primarily by R_{Edd} (Boroson & Green 1992; Sulentic et al. 2000). NLS1 galaxies were once thought to be radio-quiet. This was simply a consequence of the fact that NLS1 galaxies have a low radio-loud fraction (Zhou et al. 2006; Komossa et al. 2006b) and the small number of NLS1 galaxies known till then.

Previous studies of the radio properties of NLS1 galaxies

are all based on small samples. These showed that the radio sources, if detected, are compact (less than a few hundred parsec), at modest radio power (Ulvestad et al. 1995), of steep radio spectra (Moran et al. 2000), mostly in the RQ regime (Stepanian et al. 2003), and of low RL fraction (Zhou & Wang 2002). Until recently, there were only several RL NLS1 galaxies identified and studied individually (see Komossa et al. 2006b, for a review and references therein). Studies using (small) samples of RL NLS1 galaxies were carried out by Komossa et al. (2006b) and Whalen et al. (2006) for non-radio selected and radio selected objects, respectively. These studies confirmed that, as in ‘normal’ NLS1 galaxies (their RQ counterparts), RL NLS1 AGN are accreting at a high rate close to the Eddington limit. More importantly, as pointed out by Komossa et al. (2006b), with relatively low M_{BH} and high R , they occupy a previously sparsely populated region in the $M_{\text{BH}}-R$ diagram. Most of the sources of that sample are compact steep spectrum sources. It was confirmed that RL NLS1 galaxies are rare ($\sim 7\%$), and very radio-loud objects with $R > 100$ are even more sparse (see also Zhou & Wang 2002; Zhou et al. 2006), compared to about 10–15% for normal BLAGN and quasars (e.g. Ivezić et al. 2002). Several of these studies discussed the starburst versus AGN contribution to the radio emission, and concluded that it is AGN dominated.

Of particular interest, a few outstanding RL NLS1 galaxies came to be known to exhibit blazar properties. RXJ 16290+4007 (Schwope et al. 2000; Grupe et al. 2004) was known as a blazar and its X-ray emission is dominated by synchrotron emission (Padovani et al. 2002), or else by soft X-ray emission that is typical for NLS1 galaxies, perhaps from the accretion disk (Komossa et al. 2006b). The SED of PKS 2004-447⁶ (Oshlack et al. 2001) was well modeled with the blazar type (Gallo et al. 2006). Using information on the radio flux variations, very high radio brightness temperatures ($\gtrsim 10^{13} \text{ K}$) were inferred for J0948+0022 (Zhou et al. 2003) and 0846+513⁷ (Zhou et al. 2005)—both included in our sample, arguing for relativistic beaming (see also Wang et al. 2006). They also show some other blazar-like behavior. Recently, Doi et al. (2006, 2007) performed high-resolution VLBI observations for several radio-loud NLS1 AGN (three are included in our sample⁸) and found that they are unresolved with milli-arcsec resolutions, setting direct lower limits on the brightness temperatures in the range of $10^7 - 10^9 \text{ K}$. They also found that inverted radio spectra are common in the radio-loudest objects. The authors suggested that Doppler beaming, presumably resulting from highly relativistic jets, can explain naturally the observations. In this respect, the most remarkable object of this kind is perhaps 2MASX J0324+3410, found in our recent work (Zhou et al. 2007). It showed rapid variability in the radio, optical, and X-ray bands, and was even claimed to be marginally detected in TeV γ -rays (Falcone et al. 2004). Its non-thermal SED clearly resembles that of HBLs. Moreover, 2MASX J0324+3410 is hosted by a relatively small, apparent disk galaxy with one-armed spiral or ring galaxy morphology. These enigmatic objects present a challenge to current models of both NLS1 galaxies and blazars, and render a unique opportunity to study jet formation in black hole systems accreting at high rates.

⁶ However, the NLS1 nature of PKS 2004-447 is not certain; see Zhou et al. (2003) for a short comment.

⁷ A formerly known blazar; see Appendix B for details.

⁸ J0948+0022, J1633+4718, and J1644+2619.

⁵ The blazar sequence can be explained as, for instance, the decrease of the energy of electrons emitting at the SED peaks with increasing energy density of the seed photons for Compton scattering (Ghisellini et al. 1998)

Apparently, they do not seem to belong to any types of AGN currently known. An immediate question is whether they are some exceptional individuals or there exists a population of such objects. What are their general properties in continuum and line emission, central engine and host galaxies? What are their relations with other types of AGN?

Motivated by these issues, we compiled a RL NLS1 galaxy sample from the Sloan Digital Sky Survey (SDSS) database, aimed at a systematic study with a large and homogeneous sample. Since, as the first step, we focus on genuine RL objects, we consider in this paper objects with the radio-loudness parameter $R > 100$ (see § 2.2 for the definition of R) for the following reasons. Firstly, the classification of AGN into RL and RQ is somewhat ambiguous in the range $R = 10 - 50$ due to the presence of (a small number of) so called radio-intermediate AGN spanning in between the bulks of the two classes, which may have a different origin from genuine RL objects (e.g. Falcke et al. 2006a). The same may also be true for NLS1 galaxies. Secondly, in $R > 100$ objects contamination of radio emission from the host galaxy is negligible (see § 6.2.1). Thirdly, their jet component, if present, can be detected and studied relatively easily in other wavebands than the radio, such as optical and X-rays. Last, but not the least, they are the least studied objects *as a sample*, due to their extreme rarity, in contrast to less RL NLS1 galaxies as the bulk in the samples of Komossa et al. (2006b) and Whalen et al. (2006). We defer to a later paper a comprehensive and complete treatment of the SDSS-FIRST detected NLS1 galaxies sample, including radio-intermediate objects, to address questions such as the RL-RQ dichotomy for NLS1 galaxies. Our results show that there exists a population of NLS1 galaxies that are blazar-like, despite their rarity.

The compilation of the sample and the analyses of multi-waveband data are described in § 2. The broad band continuum and emission line properties are presented in § 3 and § 4, respectively. In § 5 the properties of their host galaxies are presented. We discuss other properties such as black hole mass and accretion rate, and isotropic radio emission, as well as the implications of the results in § 6. Throughout the paper, we assume a cosmology with $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M = 0.3$, and $\Omega_\Lambda = 0.7$. Errors are quoted at the 68% confidence level unless mentioned otherwise. We use the following conventions to denote power-law spectral indices: α_ν in the frequency domain [$S(\nu) \propto \nu^{\alpha_\nu}$, specifically $\alpha_r \equiv \alpha_\nu$ in the radio band], and α_λ in the wavelength domain [$S(\lambda) \propto \lambda^{\alpha_\lambda}$]; and the X-ray photon index Γ , defined as $f_{\text{pho}}(E) \propto E^{-\Gamma}$, where $f_{\text{pho}}(E)$ is X-ray photon number flux density.

2. SAMPLE COMPILATION AND MULTIWAVELENGTH DATA ANALYSIS

2.1. SDSS NLS1 galaxy sample

We have carried out a systematic search for NLS1 galaxies from the SDSS spectroscopic samples and the first results have been published in Zhou et al. (2006). The procedures of data analysis have been documented in that paper in detail and only an outline is summarised here. We carefully fit emission line spectra, AGN continua, and host galaxy starlight in a self-consistent manner. Firstly, galaxy starlight and AGN continuum, as well as the optical Fe II emission complex are modeled (see Appendix A for a brief account) and then subtracted. The optical Fe II multiplets are modeled with the Fe II spectral data given by Véron-Cetty et al. (2004) for both the broad and narrow components. Emission line spectra are fitted with the following models, using a code similar to that

described in Dong et al. (2005). The Balmer emission lines are de-blended into a narrow and a broad component, which are modeled by a Gaussian and a Lorentz profile, respectively. All narrow emission lines are fitted with a single Gaussian except the [O III] $\lambda\lambda 4959/5007$ doublet. Each line of the [O III] doublet is fitted with two Gaussians, one of which is used to account for a possible blue wing as seen in a few objects. As shown in Zhou et al. (2006), our analysis procedures yield reliable and accurate measurements of emission line parameters.

The SDSS spectroscopic samples of both galaxies and QSOs in the redshift range of $z < 0.8$ were analyzed based on the SDSS data release 5 (DR5, Adelman-McCarthy et al. 2007). Following Zhou et al. (2006), we classify NLS1 galaxies as those having the “broad” component of H β or H α which is detected at the 10σ or higher confidence level and is narrower than 2200 km s^{-1} in FWHM. As discussed in Zhou et al. (2006), objects selected as such fulfill naturally the second of the conventional criteria of NLS1 classification, i.e. [O III] $\lambda 5007/\text{H}\beta < 3$, and the sample is reliable and uniform. As a result, about 3300 NLS1 galaxies are found, superseding the previously published 2011 drawn from the SDSS DR3 (Zhou et al. 2006, hereafter Zhou06 sample).

2.2. Search for radio counterparts

Having compiled a large NLS1 galaxies sample from the SDSS, we search for their radio emission using data of the FIRST⁹ survey (Becker et al. 1995). The survey has a spatial resolution of $5''$ and typical 90% source positional uncertainty ellipses less than $1''$. The identification is performed by matching the SDSS positions of NLS1 AGN against those of radio sources following the procedure used by Lu et al. (2007). This procedure was designed to search for SDSS quasars detected in the FIRST survey, and is summarised here. Since extended radio sources often have multiple (diffuse) components in morphology and sometimes their positions are vague to define, we treat extended sources differently from compact sources in search of radio counterparts. By compact FIRST source we mean that only one source is found within $3'$ of the optical position and is unresolved; for them, we use a $2''$ matching radius. We search for possible extended radio sources associated with the NLS1 AGN in two steps. First, candidates are selected if they match one of the following criteria. (1) Only one resolved radio source is found within $3'$ of the optical position, and the optical position is located within the size of the radio source. (2) Two radio sources are located nearly symmetrically around the optical position within $3'$. (3) More than two radio sources are scattered around the optical position within $3'$. Then we visually inspect $6' \times 6'$ cutouts of the FIRST images centered at an object to reject false matches. As discussed in Lu et al. (2007), this approach has proved to be effective for finding extended radio sources associated with AGN; about 24% of the radio quasars in the sample of Lu et al. (2007) are resolved by the FIRST survey, among which half show FR II type morphology (Fanaroff & Riley 1974).

This procedure yields a sample of SDSS NLS1 galaxies detected in the FIRST survey. Interestingly, we note that *all* the radio sources associated with these NLS1 AGN are compact in morphology. For each object we calculate the radio-loudness parameter defined as $R_{1.4} \equiv f_\nu(1.4 \text{ GHz})/f_\nu(4400 \text{ \AA})$, where the fluxes are in the rest

⁹ Faint Images of the Radio Sky at Twenty-centimeters

frame of the objects¹⁰. The rest frame 4400 Å flux densities are calculated from the SDSS g-magnitudes assuming a spectral slope $\alpha_\nu = -0.5$. An index of -0.5 is also assumed for the K-correction for the radio fluxes. We focus on very radio-loud objects only with $R_{1.4} > 100$ in this paper, and will present the whole sample elsewhere.

We note that three NLS1 galaxies with apparent $R_{1.4} > 100$ stand out for their extremely red optical continua ($\alpha_\lambda \gtrsim 2.7$), namely, J095919.14+090659.4, J111354.66+124439.0, and J233903.82-091221.3. Our original fit to the continuum spectrum with a fixed power law slope $\alpha_\lambda = -1.7$ yields substantial reddening of $E(B-V) = 0.7-1$. Among them, J233903.82-091221.3 ($z=0.66$) has been found to show convincing evidence for substantial dust extinction (Wang et al. 2005)¹¹. Given the fact that the other two objects resemble closely J2339-0912 in their SDSS spectra, their optical light is most likely subject to heavy extinction as well. After correction for extinction, their radio-loudness parameters are largely reduced to $R_{1.4} \simeq 10-20$, below or merely close to the RL/RQ dividing line. We therefore exclude these three objects from our current sample.

2.3. Very radio-loud NLS1 galaxy Sample

The above selection procedures result in 23 NLS1 galaxies having $R_{1.4} > 100$, which compose our sample. By excluding less RL objects with $R_{1.4} < 100$ our sample can be regarded as very radio-loud to some extent. We believe that the radio-loudness estimates of our sample are not significantly affected by optical extinction, as is discussed in § 6.2.2. The objects are listed in Table 1, along with some relevant parameters of the continua and emission lines. As demonstration, we show in Figure 1 examples of the SDSS spectra and their best fit models for two representative objects; they are chosen to represent the ranges of some characterising parameters of the sample. Measurable starlight contribution is present in only one object, J1633+4718, accounting for at most $\sim 20\%$ of its observed spectral flux density at 5100 Å. Most of our objects show strong Fe II emission complexes (see Section 4). The [O III] lines are weak in general, all having $[\text{O III}]\lambda 5007/\text{H}\beta < 1.2$. All of the objects fulfill the conventional definition of NLS1 (Osterbrock & Pogge 1985; Goodrich 1989), and are therefore bona-fide NLS1 galaxies.

The redshift and radio-loudness distributions are plotted in Figure 2. Our sample has a relatively high redshift distribution peaked at $z \sim 0.5$ (the median). We calculate the B-band absolute magnitudes M_B assuming $\alpha_\lambda = -1.5$ for the K-correction, which are listed in Table 1. The M_B values range from -20.8 to -25.6 , with a median of -22.8 . Among the sample, 10 objects are as bright as $M_B \leq -23$ and hence can be classified as narrow-line type I quasars. Three objects in the sample have been previously identified with RL NLS1 galaxies and studied in detail, namely, J094857.32+002225.5 (Zhou et al. 2003), J084957.98+510829.0 (Zhou et al. 2005), and J172206.03+565451.6 (Komossa et al. 2006a). In addition, several objects were known as general RL AGN and have been studied previously, whose properties are summarised in-

dividually in Appendix B.

2.4. X-ray and UV data analysis

We have searched for X-ray emission from the sample objects using ROSAT source catalogues of the RASS and pointed observations of both the RXP (Voges et al. 1999) and WGA (White et al. 1995) catalogues. The matching radii are 5 times the given positional errors of ROSAT sources, which are typically $10''-20''$. For each match, we visually inspect their SDSS optical images to avoid spurious matches in which the X-ray source is actually associated with another nearby object within the matching radius. Of the 23 objects, 12 were detected in the RASS and 4 in pointed observations, among which 2 detected in both. Thus a total of 14 objects—more than half of the sample—were detected in X-rays with the ROSAT PSPC¹².

All of the objects except J1633+4718 do not have sufficient X-ray counts to allow proper spectral modeling. Following Schartel et al. (1996), Siebert (1996), and Yuan (1998), we estimate the X-ray photon index Γ from the two hardness ratios¹³, wherever available, assuming an absorbed power-law spectral model. Both Γ and absorption column density N_H can be estimated as free parameters, or the index alone assuming N_H to be the Galactic column density. This method had been verified to be reliable and robust under the assumption that intrinsic spectral shapes are indeed a simple absorbed power-law (Brinkmann & Siebert 1994). The X-ray fluxes in the ROSAT PSPC band (0.1–2.4 keV) are calculated from the count rates using the energy to counts conversion factor (ECF) for a power law spectrum and Galactic absorption (ROSAT AO-2 technical appendix, 1991). The ECF for each X-ray source is explicitly calculated from the ROSAT PSPC effective area¹⁴, by using information on individual Γ if obtainable, or using the mean otherwise ($\Gamma = 2.49$, see below). We list some of the information about the X-ray observations in Table 2.

For objects not detected in X-rays, we determine upper limits on their X-ray fluxes from count rate limits set by the RASS. Since the X-ray backgrounds are very low in most of the RASS observations, detection of a source is mostly determined by the source photon counts that follow Poisson statistics. We set an upper limit of source counts as 12 for an object not detected in the RASS, as often adopted (e.g. Yuan et al. 1998). The corresponding count rate limit is then calculated by using the effective exposure time at the source position as available from the RASS exposure map. Then the flux limit for each object is calculated using the same method above and assuming Γ as the mean.

Among the ROSAT detected objects, J1644+2619 was observed with Chandra with a large number of photon counts accumulated. Moreover, for J0849+5108 the hardness ratios ($HR1 = 0.97 \pm 0.11$, $HR2 = 0.28 \pm 0.15$) yield an abnormally flat spectrum $\Gamma = 0.6^{+0.36}_{-0.52}$ (see also Zhou et al. 2005). For this object we also investigate its ROSAT data in spite of the relatively low number of counts (~ 60). We perform X-ray spectral and timing analysis for J1633+4718, J1644+2619, and J0849+5108. We briefly describe here the data reduction

¹⁰ It should be noted that our definition of radio-loudness, $R_{1.4}$, is related to the previously commonly used $R_{6\text{cm}} \equiv f_\nu(6\text{cm})/f_\nu(4400\text{Å})$ (Kellermann et al. 1989) via $R_{1.4} = 1.9R_{6\text{cm}}$. Therefore, our radio-loudness cutoff corresponds to $R_{6\text{cm}} \gtrsim 50$.

¹¹ In addition to the extremely red optical continuum, its SDSS spectrum shows signatures of reddening of broad emission lines, detection of strong Ca II and Mg II absorption lines of non-stellar origin (see Wang et al. 2005, for details).

¹² Positional Sensitive Proportional Counter

¹³ Defined as $HR1 = (C_2 - C_1)/(C_2 + C_1)$, $HR2 = (C_4 - C_3)/(C_4 + C_3)$, where C_1 , C_2 , C_3 and C_4 are the number of photon counts in the 0.1–0.4 keV, 0.5–2.0 keV, 0.5–0.9 keV, and 0.9–2.0 keV bands, respectively.

¹⁴ We used those that are appropriate for the RASS or pointed observations, accordingly.




FIG. 1.— Examples of the SDSS spectra of two radio-loud NLS1 galaxies in our sample. The left-hand side panels show the rest frame spectra and the fits to the continuum with a nuclear power-law (blue color) and optical Fe II emission (magenta) model, plus host galaxy starlight contribution (green) wherever non-negligible (see § 2.1 for a description of spectral analysis). The panels on the right-hand side show a close-up of the continuum-subtracted emission line spectra and the best fits in the $H\beta$ –[O III] region. SDSS J1633+4718 (upper panel), a flat-spectrum radio source, has the lowest radio-loudness and redshift, and its spectrum is among the highest S/N; it has the largest fraction ($\sim 20\%$) of detectable host galaxy starlight contribution among the sample. SDSS J1047+4725 (lower panel), a steep-spectrum radio source, has the highest radio-loudness and redshift, and its spectral S/N is among the lowest in the sample.




FIG. 2.— Distribution of the radio loudness $R_{1.4}$ (left panel) and redshifts (right panel) of the radio-loud NLS1 galaxies sample.

procedures and present results in § 3.3. We use version 12.2 of XSPEC (Arnaud et al. 1996) for spectral modeling.

J1633+4718 (RXJ 16333+4719): The target was observed with the ROSAT PSPC-b with an exposure time of 3732 seconds on July 24th, 1993 (Obs-ID: 701549; PI: N. Bade). We retrieved the ROSAT data from the archive and use the XSELECT (version 2.3) tool of FTOOLS (Blackburn 1995) for data analysis. The X-ray spectrum is extracted from a circle of $150''$ radius, and a background spectrum is extracted from an annulus with inner/outer radii of $195/300''$. There are ~ 976 net source counts, and the count rate is 0.26 ± 0.01 ctss $^{-1}$. The spectrum in the 0.1–2.4 keV band is binned to have at least 30 counts in each bin. J1633+4718 was also detected in the RASS with a weighted exposure of 909 s, yielding ~ 185 net source counts and a count rate 0.20 ± 0.02 ctss $^{-1}$. We extract the RASS spectrum following the procedure described in Belloni et al. (1994).

J1644+2619: The object was targeted with the Chandra ACIS-S (PI: S. Laurent-Muehleisen) with an exposure time of 2946 s on June 02, 2003. The data were retrieved from the Chandra data archive and were reduced from the level-2 data set following the standard procedure using CIAO (version 3.4). There are 553 ± 44 net source counts detected, with a count rate of 0.188 ctss $^{-1}$. The 0.2–5 keV spectrum (very few counts accumulated above 5 keV) is binned to have at least 20 counts in each energy bin.

J0849+5108: The object was observed with the ROSAT PSPC-b in a pointed observation as a target in 1993, April with an exposure time of 4496 s (Obs-ID: 700821). There are only 63 net source counts detected in the 0.1–2.4 keV band. We fit the unbinned spectrum by minimizing the C-statistic (Cash 1979; Nousek & Shue 1989), which is valid for Poisson statistics appropriate to the low counts regime.

We also searched for UV data from the GALEX (Martin et al. 2005) GR2/GR3 data release¹⁵. Twelve objects in our sample have available photometric measurements in the far and/or near UV imaging bands with an effective wavelength of 1516 Å and 2267 Å, respectively. Their UV images are mostly point source like. The GALEX magnitudes (in the AB magnitude system) are corrected for Galactic extinction using $A_{FUV}/E(B-V) = 8.376$ and $A_{NUV}/E(B-V) = 8.741$, following Wyder et al. (2005).

3. BROAD BAND CONTINUUM RADIATION

3.1. GHz radio emission

3.1.1. Radio morphology and spectrum

The FIRST radio images of our RL NLS1 sources are ubiquitously unresolved at a resolution of $5.4''$, which is remarkable. This sets upper limits on the projected size of 10–40 kpc, depending on the redshift. Among our sample there are several objects observed with VLBI, including J0948+0022 (Doi et al. 2006), J1633+4718 and J1644+2619 (Doi et al. 2007), and J1505+0326 (Dallacasa et al. 1998), as reported in the literature; they remain unresolved at resolutions ranging from a few milli-arcsec to a few tens of milli-arcsec, which correspond to several to several-tens parsec in physical scale. These observations set constraints on the brightness temperatures of $T_b > 10^{8-9}$ K (Doi et al. 2006, 2007). Since all the radio sources in our sample are compact, we use hereafter the average of the FIRST and NVSS¹⁶ measurements as their




FIG. 3.— Distribution of the radio spectral indices α_r around 5 GHz [$S(\nu) \propto \nu^{\alpha_r}$].

1.4 GHz flux densities in the following analysis; this is to minimize the effects of fluctuations on the results caused by large flux variations. Similarly, at other frequencies, in cases where multiple epoch measurements are available at the same frequency, the average flux is used.

Multi-frequency and multi-epoch radio data were collected from the NASA/IPAC Extragalactic Database (NED). Of the 23 objects, 11 have flux measurements at 5 GHz, for which the 1.4 GHz–5 GHz spectral indices α_r are estimated, though the observations were not simultaneous. The 5 GHz fluxes and α_r are listed in Table 1, and the histogram of α_r is shown in Figure 3¹⁷. It can be seen that the radio spectra are systematically flat, with a median of -0.30 and a mean of -0.13 . Adopting the conventional dividing line between flat- and steep-spectrum sources, $\alpha_r = -0.5$, 8 out of the 11 objects with α_r measurement turn out to be flat-spectrum radio sources ($\alpha_r > -0.5$). In particular, several objects show an inverted radio slope ($\alpha_r > 0$) near 5 GHz and toward higher frequencies based on simultaneous multi-frequency measurements, namely, J0948+0022 (Reich et al. 2000; Zhou et al. 2003; Doi et al. 2006), J1633+4718 (Neumann et al. 1994), J1644+2619 (Doi et al. 2007), and J1505+0326 (Tinti et al. 2005). Only 3 objects have α_r steeper than, yet very close to, -0.5 .

Among the remaining 12 objects without available slopes near 5 GHz, 9 have lower frequency 327 MHz–1.4 GHz indices measurable (see Section C), out of which 6 have flat spectra ($\alpha_{327M}^{1.4G} > -0.5$). Thus, it is likely that most of these 12 objects may have the same flat (or even flatter) indices at higher frequencies as well, unless there is a break near 1 GHz, as in Gigahertz-peaked spectrum sources (GPS). This fact supports further the flat spectral index distribution found above for our sample. Though in part of the sample the radio slopes are estimated from non-simultaneous data, the current α_r distribution should be correct in the statistical sense; therefore, we expect that the bulk of our sample objects should be flat-spectrum sources. As is known, radio sources with flat spectrum, which can be produced by relativistic jet models (e.g. Blandford & Königl 1979), have a fairly good chance to be blazars (e.g. Fugmann 1988).

Now we can elaborate the radio-loudness parameter calculation for our sample by using the averaged 1.4 GHz flux densities and the estimated α_r for individual sources for the K-correction (if not available, $\alpha_r = -0.5$ is assumed as above).

¹⁷ For J0849+5108, the flux variations at 1.4 GHz were substantial and the spectral shape around 5 GHz was flat or even inverted (Arp et al. 1979); we thus use $\alpha_r = 0$ for this object in the analysis below.

¹⁵ <http://galex.stsci.edu/GR2/>

¹⁶ The NRAO VLA Sky Survey

We also make use of the measured optical slopes as presented in § 3.2. The improved radio-loudness values are listed in Table 1. The differences between the previous estimates and the improved values are small, typically within 10% in $\log R_{1.4}$; such differences are comparable with changes in $R_{1.4}$ caused by source flux variations typical of our objects (see below).

3.1.2. Radio variability and brightness temperature

A number of objects showed significant flux variations based on multi-epoch observations taken at the same wavelengths¹⁸. In Table 3 we list those having the significance of variations higher than $\sim 3\sigma$ between two observations, where σ is defined as $\sigma = (S_1 - S_2) / \sqrt{\sigma_{s1}^2 + \sigma_{s2}^2}$. We also give in the table the amplitudes of variations ΔS , the fractional amplitudes $\Delta S / \langle S \rangle$ where $\langle S \rangle$ is the average of the two fluxes, and the time spans Δt between the two observational epochs. Their variability is consistent with the fact that the variables have ubiquitously flat or inverted radio spectra when available. The fractional amplitudes given in Table 3 have a median of 43%. In particular, several objects show large amplitude variations (~ 40 –75%) on relatively short timescales, J0948+0022 (reported by Zhou et al. 2003), J0849+5108 (reported by Arp et al. 1979), and J1505+0326 within a few years, and J1548+3511 by 44% within 207 days (140 days in the source rest frame).

Large amplitude variability on short timescales has been commonly used to set lower limits on the *apparent* brightness temperature (T_B) of a radio source, since the size of the variable part of the source can be constrained from the light-crossing time (e.g. Krolik 1999; Fanti et al. 1983; Gopal-Krishna et al. 1984). Assuming that the variable portion of the radio flux is emitted from a region with a size smaller than that corresponding to the light-crossing time, which is the time span between the two observations, the *apparent* brightness temperature is estimated to be¹⁹

$$T_B \gtrsim \frac{\Delta P_{\nu e}}{2\pi^2 k \nu^2 (\Delta t)^2} = \frac{2D_L^2 \Delta S_\nu}{(1+z)\pi k \nu^2 (\Delta t)^2} \quad (1)$$

where k is the Boltzmann constant, D_L the luminosity distance, ΔS_ν the variable portion of the observed flux density, $\Delta P_{\nu e}$ the corresponding radio power at the emission frequency in the source rest frame, ν the observing frequency and Δt the time span in the observer’s frame. The estimated T_B are listed in Table 3.

Of the 8 objects, 6 have estimated T_B exceeding the equipartition brightness temperature $\sim 3 \times 10^{11}$ K as proposed by Readhead (1994), which is expected to conform by most radio sources. In particular, 4 of the objects, all of flat-spectrum, have T_B exceeding the inverse Compton limit $\simeq 10^{12}$ K (Kellermann & Pauliny-Toth 1969). The highest T_B is found to be 10^{14} K in J1505+0326, from two VLBI observations at 8.3 GHz with the variability significance of 3σ . The

extremely high T_B values are commonly explained as emission originating from relativistic jets (Blandford & Königl 1979; Jones & Burbidge 1973). For instance, taking the inverse Compton limit as a conservative limit, the minimum Doppler factor can be estimated as $\delta_{\min} = (T_B / 10^{12} \text{ K})^{1/3}$. This results in $\delta_{\min} = 1.5$ –4.7 for the 4 objects (Table 3), consistent with the range of the Doppler factors of relativistic jets inferred for classical RL AGN (Ghisellini et al. 1993; Jiang et al. 1998).

3.1.3. Summary of the GHz radio properties

(1) All the radio sources of our sample are compact, being unresolved at the FIRST resolution of several arc-seconds. (2) The majority of objects with available radio indices are of flat-spectrum (around 1.4–5 GHz), some even with inverted spectra. (3) Most of the flat-spectrum sources show significant variability, from which the brightness temperatures are inferred to be as high as $T_B > 10^{11}$ K, and even exceeding the inverse Compton limit of 10^{12} K in 4 objects. These high T_B values may suggest the presence of at least mildly relativistic beaming of the radio emission.

The properties of the radio emission at low frequencies of our sample objects are summarised in Appendix C, which is important for understanding the intrinsic radio power and radio-loudness of these objects, as discussed in Section 6.3.

3.2. Optical continuum emission

Our standard algorithm to select NLS1 galaxies as above does not give information about the optical continuum shape. However, for RL objects the optical continuum may deviate from the ‘canonical’ shape in consideration of a potential contribution of jet emission. To properly take this effect into account, we parameterise the optical continuum shape especially for our RL sample. We model the AGN continuum in the 2500Å–5500Å rest frame range with a power-law or a broken power-law, and repeat the above spectral fitting with the slope as a free parameter. The fits are performed in the same way as described in § 2.1 assuming no intrinsic reddening. The improved fits give a more realistic description of the AGN continuum; while, on the other hand, we find that the new fits have almost no effects on the parameterisation of the emission line spectra. The optical continuum fluxes in the source rest frame are measured directly from the power-law fits of the AGN continuum.

It has been established that the optical luminosity L_{opt} of NLS1 AGN is tightly correlated with the H β line luminosity $L_{\text{H}\beta}$ based on large samples (Zhou et al. 2006); hence $L_{\text{H}\beta}$ is a good indicator of L_{opt} . We test this scaling relation for RL NLS1 galaxies as shown in Figure 4, where the measured 5100Å luminosities are plotted versus $L_{\text{H}\beta}$. As a comparison the ~ 2000 predominantly RQ NLS1 galaxies in the Zhou’06 sample are over-plotted, along with the fitted $\lambda L_{\lambda 5100} - L_{\text{H}\beta}$ relation (Zhou et al. 2006, their Eq. 5). As can be seen, the RL NLS1 galaxies reveal the same trend of correlation, but, of particular interest, lie systematically above the known $L_{\text{opt}} - L_{\text{H}\beta}$ relation for normal NLS1 galaxies. We estimate the ratio of the observed luminosity to that expected from the H β luminosity ($L_{\lambda 5100} / L_{\lambda 5100}^{\text{H}\beta}$) using the $\lambda L_{\lambda 5100} - L_{\text{H}\beta}$ relation. We find that the ratios range from 0.93 to 5.2, with a mean of 1.9. The ratios differ systematically between flat- and steep-spectrum sources as groups, with the mean being 2.4 for the former and 1.8 for the latter, and 1.6 for those without radio indices.

¹⁸ In cases where two observations have much different spatial resolutions, we consider the variability to be genuine only if the higher flux was measured at a higher resolution. This is to avoid detection of spurious variations caused by the possibility that a measured higher flux may be contributed by contaminating extended/nearby emission if the observation is made at a lower spatial resolution. We take this approach even though this problem actually has little effect to our objects, all of which are compact on the scales concerned, e.g. several arc-seconds. Thus, our results are conservative regarding the number of variables, and variability amplitudes and timescales.

¹⁹ We noted that there was an error in our previous calculation of T_B in Wang et al. (2006) and Zhou et al. (2003), which was also pointed out by Ghosh & Punshi (2007). We have corrected it in this paper.




FIG. 4.— Relationship between the nuclear monochromatic luminosity at 5100Å and the H β luminosity for the radio-loud NLS1 AGN of our sample (filled dots: flat-spectrum radio sources; open circles: steep-spectrum radio sources; asterisks: no available radio indices). Over-plotted are NLS1 galaxies in the Zhou’06 sample (small dots), predominantly radio-quiet, as well as the best-fit relation (dashed). It shows that the radio-loud NLS1 galaxies have systematically higher observed luminosities in excess of that predicted from the H β luminosity of normal NLS1 galaxies.

The optical continua in the rest frame 2500Å–5500Å range can be described with a power-law for most of the objects. The fits are relatively poor for two objects, namely, J1037+0036 and J1634+4809, which show convex-shaped spectra in the optical–UV band with a drop short-ward of Mg II λ 2800Å in wavelength. For one object, J1138+3653, the spectrum cannot be fitted with a power-law due to a rapid drop in the blue part, which results possibly from mild extinction and is discussed further in §6.2.2. The best-fit slopes show a large range from modestly red ($\alpha_\lambda = -0.2$) to blue ($\alpha_\lambda = -2.4$) spectra. Their distribution is shown in Figure 5 (excluding J1138+3653), which has a median of -1.54 and a mean of -1.43 . We compare this α_λ distribution with that of a RQ NLS1 sample, which comprises 55 SDSS-selected objects with reliably measured optical slopes (dashed line in the figure, Ai et al. 2008, for objects with multi-epoch data the averaged slopes are used). It should be noted that that RQ sample has the redshift and luminosity distributions indistinguishable from those of our sample [the two-sided Kolmogorov-Smirnov (K-S) test probability of 0.48 and 0.52, respectively]. It can be seen that, though there is a broad range of overlap, our very radio-loud NLS1 AGN appear to have systematically bluer continua than the RQ objects. The RQ sample have a median of -1.24 and a mean of -1.15 . The chance probability that the two distributions are the same is only $P_{\text{chance}} \lesssim 0.01$, as given by the K-S test. A comparison with the smaller NLS1 galaxies sample of Constantin & Shields (2003), which have UV–optical slope measurements with the Hubble Space Telescope (HST), yields a similar conclusion; the closest value of their sample to ours is the continuum slope of the median composite spectra, $\alpha_\nu \sim -0.79$ ($\alpha_\lambda \sim -1.21$).

3.3. X-ray radiation




FIG. 5.— Histogram of the power-law slopes of the optical continuum (rest frame 2500–5000Å) for our radio-loud NLS1 sample (solid line; except for one object which is fitted with a broken power-law). As a comparison, the dashed line represents the histogram of α_λ for a radio-quiet NLS1 sample selected from the SDSS with redshift and luminosity distributions compatible to ours (from Ai et al. 2008).

Assuming an absorbed power-law spectral model and using the aforementioned hardness ratio method, the effective photon indices are estimated for most of the objects with valid ROSAT/PSPC hardness ratios extracted from the ROSAT source catalogues. The estimated absorption N_H values are found to be well consistent with the Galactic values within errors, indicating that there is no significant X-ray absorption in these objects. Hence, we adopt the photon indices Γ (Table 2) obtained by assuming $N_H = N_H^{\text{Gal}}$ for better constraining the parameters. Furthermore, for 5 of the objects no physically meaningful result can be obtained by this method, because either the spectrum is complex and highly deviating from the assumed absorbed power-law, or the measured hardness ratios are displaced outside the valid zone by Poisson fluctuations. For objects with multiple X-ray data, we use the weighted mean for Γ and fluxes in the following analyses. Results from detailed spectral analysis for the 3 objects, J1633+4718, J1644+2619, and J0849+5108, are summarised below.

J1633+4718 (RX J16333+4719): The spectrum deviates apparently from a simple absorbed power-law model, with prominent soft X-ray emission dominating $E < 0.5$ keV and a hard tail at $E > 0.5$ keV. The best fit is achieved using a model of a power-law plus a redshifted blackbody component, and the absorption N_H is found to be in excellent agreement with the Galactic $N_H^{\text{Gal}} = 1.79 \times 10^{20} \text{ cm}^{-2}$. Hence N_H is fixed to N_H^{Gal} . The data and the best fit model are shown in Figure 6. The rest frame blackbody temperature is found to be $32.5^{+8.0}_{-6.0}$ eV and the photon index of the underlying power law $\Gamma = 1.37 \pm 0.49$ (90% confidence for one parameter of interest). The latter is less well constrained due to the limited bandpass of the PSPC in the ‘hard’ X-rays. No temporal variability is found within the observation interval. The spectrum measured in the RASS has almost identical spectral shape as that of the pointed observation. The best-fit parameters are in excellent agreement with those obtained from the pointed observation, with $\Gamma = 1.47^{+0.77}_{-0.92}$ and the blackbody tempera-

ture 30^{+12}_{-10} eV (90% confidence level), though the errors are large given the small source counts. No variation in either the spectral shape or the flux is found between the RASS and the pointed observation.

J1644+2619: A simple power-law model is found to be an excellent fit to the spectrum (see Figure 7), resulting in a reduced $\chi^2 = 13$ (for 23 d.o.f). The fitted absorption N_{H} is almost identical to the Galactic value $N_{\text{H}}^{\text{Gal}} = 5.12 \times 10^{20} \text{ cm}^{-2}$. The best-fit photon index is steep, $\Gamma = 2.19 \pm 0.27$ (90% confidence for one parameter of interest). Adding an extra soft X-ray component (e.g. blackbody) does not improve the fit, though with two more free parameters; the power-law photon index becomes flatter, $\Gamma \sim 1.8^{+0.6}_{-0.3}$, but is subject to a large uncertainty range which encompasses the single power-law model. We therefore suggest that the simple power-law model with a relatively steep index is the most likely spectral shape, though models composed of a flatter power-law with a weak soft component cannot be ruled out from the current data. Compared to the observation in the RASS, the X-ray source was about 2.5 times fainter at the Chandra observation epoch, though the photon index seems not to vary.

J0849+5108: The spectrum can be fitted with a power-law with absorption N_{H} close to the Galactic value $N_{\text{H}}^{\text{Gal}} = 3.0 \times 10^{20} \text{ cm}^{-2}$. Fixing column density at $N_{\text{H}}^{\text{Gal}}$ yields a photon index $\Gamma = 1.77^{+0.44}_{-0.60}$ (90% confidence for one parameter of interest). We note that the extremely hard hardness ratios given in the source catalogue are incorrect, which may result from inappropriate background subtraction around this object in the presence of nearby, seemingly extended X-ray emission.

Figure 8 shows the distribution of the *effective* X-ray photon indices Γ , which span a wide range from 1.7 to 3.3. Since the uncertainties of most of the derived Γ are relatively large, we try to disentangle the intrinsic distribution of Γ from the measurement uncertainties, using the Maximum-Likelihood method introduced by Maccacaro et al. (1988). Assuming that the intrinsic distribution of Γ is Gaussian, we find that the distribution is intrinsically broad, with a standard deviation of $0.39^{+0.41}_{-0.19}$ and a mean of $2.60^{+0.37}_{-0.43}$ (at the 90% confidence level for two interesting parameters). We conclude that the soft X-ray continuum shape, as described by the power-law photon index Γ , exhibits a considerable variety among our RL NLS1 galaxies, with both flat and steep spectra.

It should be noted that the thus estimated Γ from hardness ratios are only an indicator of the overall spectral shape in the 0.1–2.4 keV range. A steep Γ value does not rule out the presence of a flat ‘hard’ X-ray component if both a steep soft X-ray and a flat ‘hard’ X-ray component co-exist. As an example, for J1633+4718, a slope of $\Gamma \simeq 2.8$ is derived from its hardness ratios if the spectrum is approximated as a power-law. This is particularly true for NLS1 galaxies, since a soft X-ray excess component is commonly seen in them. Better quality X-ray spectra in a wider energy band than that of the ROSAT/PSPC are needed to determine the true underlying power-law in these objects.

3.4. Broad band SED

To quantify the broad band SED, we calculate the commonly used effective spectral indices between the 5 GHz radio, 2500 Å optical/UV, and 2 keV X-ray band, namely, α_{r0} , α_{ox} , and α_{rx} ; they are defined as $\alpha_{12} = -\log(S_{\nu 1}/S_{\nu 2})/\log(\nu_1/\nu_2)$, where $S_{\nu 1}$ and $S_{\nu 2}$ are the fluxes at the frequencies ν_1 and ν_2 , respectively, in the object rest

frame²⁰. The rest frame 2500 Å fluxes are computed from the *u*-band PSF magnitudes, which have the effective wavelength (3542 Å) very close to the redshifted 2500 Å light (within a range of 500 Å); for the K-correction, the measured optical slopes for individual objects are adopted (for one object without the slope available, $\alpha_{\lambda} = -1.5$ is assumed). For the 3 objects for which X-ray spectral fits were enabled, the X-ray flux densities at 2 keV ($f_{2\text{keV}}$) are measured directly from the best-fit spectral model; while for the others $f_{2\text{keV}}$ are computed from the 0.1–2.4 keV integrated fluxes. In doing so, as well as the K-correction, the derived Γ values are used for individual sources whenever available, or the mean $\langle \Gamma \rangle = 2.6$ is used otherwise. For those without X-ray detection, lower limits on α_{ox} and α_{rx} are calculated using the upper limits on the X-ray fluxes constrained by the RASS. The obtained α_{r0} , α_{rx} , and α_{ox} are listed in Table 4.

In Figure 9 we plot our RL NLS1 galaxies on the commonly used $\alpha_{\text{r0}}-\alpha_{\text{ox}}$ blazar diagnostic diagram, separating them into three groups: flat (underlying) spectra ($\Gamma < 2.0$, filled dots), steep spectra ($\Gamma > 2.0$, open circles), and those without Γ estimates (asterisks). For comparisons, we over-plot RLQs from the ROSAT detected quasar sample of Brinkmann et al. (1997). That sample is composed of both FSRQ and steep-spectrum radio quasars (SSRQ), whose loci largely overlap with each other. It can be seen that some of the objects do follow the trend defined by FSRQ/SSRQs, indicating their SEDs are similar to those of FSRQ/SSRQs. Specifically, all the 3 objects in our sample known to have a steep radio spectrum are well consistent with normal SSRQs. It is known that FSRQs (and SSRQs) occupy almost the same region on the $\alpha_{\text{r0}}-\alpha_{\text{ox}}$ diagram as LBLs, owing to their similar SEDs (Padovani et al. 1997a; Brinkmann et al. 1996; Sambruna et al. 1996; Fossati et al. 1998).

However, a considerable fraction of the objects clearly offsets. We expect that these outliers must have different SEDs from those of classical FSRQ and LBLs. By invoking the classification scheme of BL Lac objects, we divide the $\alpha_{\text{r0}}-\alpha_{\text{ox}}$ diagram into the loci of LBLs and HBLs. A simple yet commonly used criterion separating HBLs and LBLs is α_{rx} ; a nominal value is set to be around $\alpha'_{\text{rx}} \sim 0.75 - 0.78$ (defined between 5 GHz and 1 keV) suggested by various authors, with HBLs having smaller (flatter) α_{rx} values. We adopt 0.78 as the dividing line (as in, e.g. Padovani et al. 2003), which corresponds to $\alpha_{\text{rx}} = 0.787$ for the α_{rx} definition adopted in this paper and is represented by the dashed line in Figure 9. Following Padovani et al. (2003), a more elaborated locus for HBL is also indicated schematically, which is populated by X-ray selected BL Lac objects (e.g. Brinkmann et al. 1996). Of particular interest, most of the objects offsetting from the classical FSRQ and LBL locus actually fall within the HBL locus²¹. This indicates that the broad band SEDs of these NLS1 AGN are similar to those of HBLs. Follow-

²⁰ There are slight differences in the frequencies adopted in the α_{r0} and α_{ox} definitions among the literature. Whenever we cite α_{r0} and α_{ox} which have different definitions, we first convert their values to those according to the definition used in this paper.

²¹ In a strict sense, the HBL locus in Figure 9 represent the SEDs of the ‘pure blazar’ component only or dominated by the non-thermal continuum. Our NLS1 objects are, however, strong-lined and a significant contribution is expected from the thermal ionizing continuum to the observed optical flux ($\sim 50\%$ on average, see § 3.2). For rigorous comparisons with HBLs, we should consider the ‘corrected SED’ corresponding to the pure non-thermal continuum only. However, such corrections are very small ($\Delta\alpha_{\text{ox}} \sim -0.12$ and $\Delta\alpha_{\text{r0}} \sim 0.05$ on average) for the bulk of the objects, and the results are essentially not affected.




FIG. 6.— X-ray spectral fit for J1633+4718. Left panel: ROSAT PSPC spectrum, the folded best-fit model and the residuals; right panel: best fit model of a power-law plus a black-body (with Galactic absorption).




FIG. 7.— Chandra ACIS X-ray spectrum of J1644+2619 and the best-fit model as a steep power-law ($\Gamma = 2.19 \pm 0.27$, 90% errors) with Galactic absorption.

ing our terminology introduced in § 1, we refer to them as NLS1 HFSRQs, and those with SEDs typical of classical FSRQs (LBL-like) as NLS1 LFSRQs. These NLS1 HFSRQs have ubiquitously flat radio spectra whenever available, consistent with a potential blazar nature. Those without X-ray detection (arrows) have their α_{ro} and α_{ox} limits consistent with the α_{ro} and α_{ox} values for either LFSRQs/SSRQs or HFSRQs. Also, a few objects lie very close to the border line, on both the HBL and the LBL sides, suggesting a smooth transition between the two types. As a comparison, we also mark in the plot the position of 2MASX J0324+3410, an extreme NLS1–blazar composite object, which has a HBL-type SED with the synchrotron peak in the UV/soft X-rays (Zhou et al. 2007). Clearly, 2MASX J0324+3410 fulfills the HBL classification in terms of its radio–optical–X-ray SED, as expected. We note there are a few objects lying in the vicinity of 2MASX J0324+3410 in the $\alpha_{\text{ro}}-\alpha_{\text{ox}}$ diagram, suggesting their similarities and thus the previously known NLS1–blazar hybrid 2MASX J0324+3410 is not unique.

FSRQs/LBLs and HBLs are known to differ in their X-ray spectral shapes as FSRQs/LBLs have relatively flat spectra




FIG. 8.— Distribution of the *effective* photon indices of representing power-law of the X-ray spectra of our radio-loud NLS1 galaxies, where available. All the indices are measured with the ROSAT PSPC in 0.1–2.4 keV, except for J1644+2619 with the Chandra ACIS in 0.2–5 keV. For J1633+4718 we adopt $\Gamma = 2.8$ obtained from the hardness ratios to represent the overall spectral shape.

while HBLs have steep spectra (e.g. Worrall & Wilkes 1990; Brinkmann et al. 1996; Padovani & Giommi 1995; Wolter et al. 1998); specifically, $\Gamma \sim 2.0$ were found for FSRQs/LBLs and $\Gamma \sim 2.5$ or steeper for HBLs in the ROSAT band. As such, the overall spectral shapes from the optical to the X-ray band are concave in FSRQs/LBLs ($\alpha_{\text{ox}} > \alpha_x$, where $\alpha_x \equiv \Gamma - 1$ is the energy index), while they are convex ($\alpha_{\text{ox}} < \alpha_x$) in HBLs (e.g. Sambruna et al. 1996; Padovani et al. 1997a; Perlman et al. 2005). Parallely, the X-ray emission of HFSRQs is hypothesized to be dominated by synchrotron radiation with a steep X-ray spectrum, as claimed to be seen in a few individuals (Padovani et al. 2002). For the NLS1 HFSRQs in our sample with measured X-ray slopes, we find that all but one (J1633+4718) have steep X-ray spectra ($\Gamma > 2.0$). We also examine the optical-to-X-ray spectral shapes for the objects with available α_x by plotting α_x versus α_{ox} in Figure 10. As can be seen, the two objects (J1633+4718 and J0849+5108) with flat X-ray spectra $\alpha_x < 1.0$ show prominently concave optical-to-X-ray spectral shapes; whereas most of the steep X-ray spectrum objects are consistent with convex spectra, including 2MASX J0324+3410 with $\Gamma \simeq 2.2$ (Zhou et al. 2007).




FIG. 9.— α_{ro} versus α_{ox} for the radio-loud NLS1 sample; filled dots: objects with flat X-ray spectra ($\Gamma < 2$); open circles: objects with steep X-ray spectra ($\Gamma > 2$); asterisks: objects without available X-ray slopes. Arrows represent lower limits of α_{ox} for X-ray non-detections. For comparison, we over-plot ROSAT detected radio-loud quasars from Brinkmann et al. (1997), consisting of both FSRQs (small crosses) and SSRQs (small dots). The blazar-NLS1 hybrid object 2MASX J0324+3410 discovered by Zhou et al. (2007) is also indicated for comparison (square). The dashed line represents a constant $\alpha_{rx} = 0.78$, which is commonly used to divide BL Lac objects into LBLs and HBLs (see text for details). A more conservative, schematic locus for HBL is also indicated following Padovani et al. (2003, solid line).




FIG. 10.— X-ray spectral energy index α_X ($\Gamma - 1$) versus α_{ox} . The overall optical-to-X-ray spectral shapes are divided into convex shapes ($\alpha_{ox} < \alpha_X$) and concave shapes ($\alpha_{ox} > \alpha_X$) by the dashed line ($\alpha_{ox} = \alpha_X$). The numbers mark the object IDs as given in Table 1. 2MASX J0324+3410 is also plotted (square).

We conclude that some of our radio-loud NLS1 AGN have SEDs similar to those of LBLs/RLQs, while the remaining similar to HBLs/HFSRQs. The X-ray spectral shapes are also broadly consistent with these two types of SED.

Finally, in Figure 11 we plot the broad band SED in the $\log \nu - \log \nu L_\nu$ representation for the 14 objects with X-ray detection. Though there is only one broad radiation hump

detected (due to a lack of hard X-ray and γ -ray data), they do reveal some similarities with the low-energy (synchrotron) hump characteristic of blazars. In particular, some of the HBL-like candidates are found to have the radiation hump peaked around the UV, supporting the results from the above $\alpha_{ro}-\alpha_{ox}$ distribution; these include J1146+3236, J1238+3942, J1644+2619, and J1722+5654, etc. They also show SEDs resembling that of 2MASX J0324+3410.

4. EMISSION LINES AND AGN EIGENVECTORS

The Eigenvector-1 space involves primarily a set of well known correlations among the following parameters: the line width $FWHM(H\beta)$, R_{4570} , $[O III]\lambda 5007/H\beta$, and soft X-ray spectral index (Boroson & Green 1992). R_{4570} is the relative strength of the Fe II multiplets in terms of the Fe II-to- $H\beta$ flux ratio, $R_{4570} \equiv Fe II/H\beta$, where Fe II is the flux of the Fe II multiplets in the range of 4434–4684 Å, and $H\beta$ is the total $H\beta$ flux (Véron-Cetty et al. 2001). $[O III]\lambda 5007/H\beta$ is the strength of $[O III]$ relative to that of $H\beta$. Here we examine these correlations for our RL NLS1 galaxies, and compare them with those for the large NLS1 sample of Zhou'06, which are predominantly radio-quiet (plotted as small dots in the figures in this section).

Figure 12 shows the correlation between the $H\beta$ line-width and the relative strength of Fe II; again, a significant anti-correlation is present for the RQ NLS1 sample (with a large scatter though), as shown in Zhou et al. (2006), whereas no significant correlation is found for the RL objects. A similar result is found for the correlation between R_{4570} and $[O III]\lambda 5007/H\beta$, as shown in Figure 13. While an anti-correlation between the Fe II and $[O III]$ strength is evident for 'normal' NLS1 galaxies, there appears no such correlation for the RL objects. However, it should be noted that, given the weakness of most of these correlations found for the RQ sample, our RL NLS1 sample is too small to draw statistically significant inferences. Also, we note that the high R_{4570} values for the RL objects span a relatively narrow range of R_{4570} in the diagram, which may partly account for the lack of the $R_{4570}-[O III]\lambda 5007/H\beta$ correlation.

Another well known Eigenvector-1 correlation is the anti-correlation between soft X-ray photon index Γ and $H\beta$ line-width (e.g. Wang et al. 1996; Boller et al. 1996), that was found to extend down to $FWHM \sim 1000 \text{ km s}^{-1}$ (Zhou et al. 2006, their Figure 17). We examine this property for the 14 RL NLS1 galaxies with measured Γ , and find that they do not follow the $\Gamma-FWHM$ trend defined by their RQ counterparts. However, the sample size is small, and the scatter in the known relation is huge. Alternatively, the finding could be attributed to a different X-ray radiation mechanism dominating the ROSAT PSPC band in RL and RQ NLS1 galaxies (see below).

We also search for potential correlations between radio-loudness $R_{1.4}$ and R_{4570} as well as $[O III]\lambda 5007/H\beta$ within our RL NLS1 sample; no significant correlations are found, however.

In the process of optical spectral analysis, we have noticed that the Fe II multiplets are generally strong in our RL NLS1 galaxies. We plot in Figure 14 the distribution of R_{4570} , and, as a comparison, also the R_{4570} distribution for the NLS1 galaxies in the Zhou'06 sample (normalized to the number of RL objects). The two-sided K-S test testing if the two distributions are drawn from the same population yields a probability level of 0.2% only; it is evident that the RL NLS1 galaxies have systematically higher R_{4570} val-




FIG. 11.— Broad-band spectral energy distribution in the $\log \nu L_\nu - \log \nu$ representation for radio-loud NLS1 AGN with available X-ray data. The luminosities are at the emitting frequencies in the rest frame of the objects. The radio and infrared data are collected from the NED and supplementarily from recent radio surveys such as FIRST. The optical data are SDSS PSF magnitudes in the five bands (u, g, r, i, z). The UV data are GALEX near- and far-UV magnitudes. The optical and UV magnitudes have been corrected for Galactic extinction. The host galaxy contamination should be negligible, as shown in Section 2.3. For objects whose X-ray spectrum can be represented by a power-law, the estimated slope and uncertainties are indicated as a bow-tie. For J1633+4718, the X-ray spectrum is complex and the unfolded spectrum is presented.

ues than their RQ counterparts. The average R_{4570} is 1.03, compared to 0.82 for the ‘normal’ NLS1 galaxies with reliable Fe II measurements in the Zhou’06 sample. To examine whether the large R_{4570} values are caused by systematically weak (total) $H\beta$ emission, we compare the $H\beta$ flux distribution of the RL and RQ samples; we find that the two $H\beta$ flux distributions are indistinguishable between the two samples (the K-S test). Moreover, the RL NLS1 sample has $FWHM(H\beta)$ and $[O\text{ III}]\lambda 5007/H\beta$ distributions indistinguishable from those of the Zhou’06 sample (the K-S test). This result, together with the absence of the anti-correlations of $R_{4570}-FWHM(H\beta)$ and $R_{4570}-[O\text{ III}]\lambda 5007/H\beta$ indicate that the high R_{4570} values for the RL sample are real, rather than spuriously induced from dependence on other parameters. We thus conclude that the optical Fe II emission in our very RL NLS1 galaxies is on average stronger than that in RQ NLS1 galaxies. In fact, this effect has been noted in the sam-

ple of Komossa et al. (2006b, their Figure 3), but the statistical significance is not as high due to a factor of 2 smaller of the sample size.

We have identified four [O III] blue outliers in our sample, with the [O III] lines significantly blueshifted by $V([O\text{ III}]) > 200 \text{ km s}^{-1}$ relative to [O II]²². The phenomenon of blue outliers has been observed in a few radio galaxies (e.g. Tadhunter et al. 2001), and is especially frequent among NLS1 galaxies (Komossa et al. 2007, and references therein). We find a fraction of blue outliers with $V([O\text{ III}]) > 200 \text{ km s}^{-1}$ of 17% among our sample, about a factor of 2.4 higher than that in the RQ NLS1 sample of Komossa et al. (2007). These authors also discovered that, for RQ blue outliers, the velocity offsets are correlated with ionization potential and line width. Such correlations persist for RL blue outliers. In all four

²² $V([O\text{ III}])$ are 220, 400, 480 and 520 km s^{-1} for J1634+4809, J1505+0326, J1305+5116, and J1443+4725, respectively




FIG. 12.— Ratio of the Fe II multiplets to the total $H\beta$ line flux, R_{4570} versus $H\beta$ line-width in $FWHM$ for the radio-loud NLS1 galaxies (fill circles), as well as normal NLS1 galaxies in the Zhou et al. (2006) sample (small dots).




FIG. 13.— Ratio of the Fe II multiplets to the total $H\beta$ flux, R_{4570} versus the ratio of the [O III]-to- $H\beta$ line flux. Plot symbols are the same as in Figure 12.

we detect blueshifted [Ne V] lines with velocity shifts up to $\sim 2300 \text{ km s}^{-1}$ (J1443+4725). The high frequency of blue outliers in RL NLS1 galaxies may indicate that jet-cloud interactions contribute to, or dominate, the [O III] $\lambda 5007$ blueshifts in the RL sources.

Most of the objects show symmetric profiles of the [O III] $\lambda 5007$ line. In three objects, particularly J1633+4718, the [O III] doublets show a blue wing that can be modeled by an extra Gaussian component (see Figure 1). This indicates the presence of possible outflows associated with the narrow line region. In addition, J103123.73+423439.3 shows double-peaked profiles in both [O III] $\lambda 5007$ and $H\beta$.

5. HOST GALAXIES

The host galaxies of radio-loud NLS1 AGN are poorly understood due to very sparse observations. The only ob-




FIG. 14.— Distribution of $Fe II/H\beta$ for the radio-loud NLS1 galaxies (solid line). For comparison, also plotted is the distribution for the normal NLS1 galaxies in the Zhou et al. (2006) sample, normalized to the size of the radio-loud sample (dashed line).

ject, as far as we are aware of, observed with the HST is 2MASX J0324+3410. Its HST image reveals a ring or one-armed galaxy morphology, indicating a possible galaxy interacting/merging history (Zhou et al. 2007). As for our current sample, given the relatively high redshifts, most of their host galaxies cannot be examined with the SDSS images. The exceptions are J1633+4718 and J1644+2619, which have redshifts of 0.116 and 0.144, respectively. Interestingly, the host galaxy of J1633+4718 has a size of more than 20 kpc along the major axis and is in an interacting pair/merger system, in which the second galaxy hosts a reddened starburst nucleus (Bade et al. 1995). Both of the member galaxies appear to be gas rich with disturbed morphology, and thus of late morphological type. The host galaxy of J1644+2619, though being compact (~ 15 kpc), also seems to be disk-like. We note that the black hole masses in both galaxies are small, 2.0×10^6 and $8.4 \times 10^6 M_\odot$ for J1633+4718 and J1644+2619, respectively (see § 6.4).

The SDSS spectrum of J1633+4718 includes a significant contribution of the host galaxy light within the fiber of $3''$ diameter, which appears to be dominated by a young stellar population, as can be seen in Figure 1 (the contamination from the starburst nucleus of $4''$ away should be negligible). Besides, marginal imprints of high-order Balmer absorption lines, though being weak, are likely present in the SDSS spectra of other three objects at higher redshifts (J1146+3236, J1505+0326, and J1634+4809; $z = 0.4 - 0.5$), indicating possible young stellar populations in their host galaxies.

We thus suggest that some of the radio-loud NLS1 AGN (at least those with a small black hole mass; see § 6.4) may reside in gas-rich, disk galaxies, and some reside in galaxies with possible young stellar populations. This result contrasts clearly with classical RLQs and BL Lac objects, that preferentially reside in giant elliptical or massive bulge-dominated galaxies, or strongly passively evolving galaxies (e.g. Bahcall et al. 1997; Dunlop et al. 2003; Floyd et al. 2004; O'Dowd & Urry 2005). While it is, to some extent, in line with the finding of Ho & Peng (2001) and Ho (2002) for the Seyfert galaxies in their samples, which are much less luminous and

more sub-Eddington than our radio-loud NLS1 AGN studied here, however. As is shown in § 6.4, most of the objects in our sample have relatively small black hole masses ($M_{\text{BH}} \sim 10^{6-8} M_{\odot}$). Hence, considering the M_{BH} -spheroid mass relation known for nearby galaxies (Magorrian et al. 1998), our finding about the host galaxies of radio-loud NLS1 AGN is not unexpected. Deeper observations are needed to reveal the host galaxy properties for the remaining radio-loud NLS1 AGN with larger M_{BH} , which are not an easy task given their relatively high redshifts, however.

6. INTERPRETATION AND DISCUSSION

6.1. The sample

Our objects fulfill the conventional definition of NLS1 galaxies, and show typical NLS1 characteristics such as strong optical Fe II emission. They also share similar distributions in black hole mass and Eddington ratio with ‘normal’ NLS1 galaxies (see § 6.4). Therefore, they should be bona-fide NLS1 galaxies. The 5 GHz radio luminosities $L_{5\text{GHz}}$ range from $10^{24.17}$ to $10^{26.96} \text{ W Hz}^{-1}$ with a median of $10^{25.23}$. Thus, most of our objects would remain qualified as radio-loud even if the alternative criterion of RL AGN is applied, which is based solely on radio luminosity, $L_{5\text{GHz}} \gtrsim 10^{25} \text{ W Hz}^{-1}$ as proposed by some authors (e.g. Miller et al. 1990; Kellermann et al. 1994). Of our sample, 19 were spectroscopically targeted in the SDSS as quasar candidates based on their optical colors, and 3 were targeted serendipitously; only one object was selected solely as a ROSAT/FIRST counterpart. Hence our very RL NLS1 sample can be regarded as an essentially optically-selected one, similar to other SDSS radio-quiet NLS1 samples.

The sample spans a luminosity range from Seyfert galaxies to quasars, with roughly half of the sample in each category. Compared to the ‘normal’ NLS1 galaxies in the Zhou’06 sample, which have an average $H\beta$ luminosity $\langle L_{H\beta} \rangle = 3.0 \times 10^{42} \text{ erg s}^{-1}$, our RL objects have relatively higher luminosities ($\langle L_{H\beta} \rangle = 6.0 \times 10^{42} \text{ erg s}^{-1}$, see also Figure 4). This difference is likely a consequence of the trend that RL objects are more likely to be found in AGN with more massive black holes (and thus higher luminosities), as suggested recently (e.g. Best et al. 2005). We find that such very RL objects represent only 0.7% of NLS1 galaxies drawn from the SDSS, though this fraction may be somewhat uncertain given all kinds of selection effects introduced in the surveys.

Compared to the other two previously published RL NLS1 samples of Komossa et al. (2006b) and Whalen et al. (2006), which comprise 11 and 16 objects respectively, our current sample differs mainly in that it has a much higher radio-loudness cutoff. Consequently, the sample is biased for selecting flat-spectrum sources dominated by beamed radio emission. In addition, our sample has a statistically higher redshift distribution than those two samples. In comparison, only three and four objects from those two samples satisfy $R_{1.4} > 100$, respectively, among which, two from the former and one from the latter sample are included in our current sample.

6.2. Broad band continuum emission

6.2.1. Radio emission: relativistic beaming and jets

The radio luminosities of our sample objects are well above those expected from the most radio-luminous starburst galaxies $\sim 10^{22.3-23.4} \text{ W Hz}^{-1}$ (Smith et al. 1998). In addition, no imprints of vigorous starbursts are found in the optical SDSS

spectra of sources of our sample; these spectra sample the central region of the host galaxies within a radius of 3–11 kpc. None of the objects are found to be associated with IRAS sources except for J1633+4718, for which the infrared emission is expected (at least predominantly) from the starburst nucleus 4'' away, rather than from the active nucleus, however. These results suggest that for essentially *all* of the objects the radio emission is of AGN origin, rather than of starburst origin.

Large amplitude flux variations up to $\sim 70\%$ have been observed in some of our objects with flat radio spectra. To make sure that this is not spurious due to contamination of nearby sources, we visually examined the FIRST and NVSS images of the variable objects. We find that for 6 of the 8 objects no other sources are detected within 3'; for the remaining 2, faint sources are present but are all well separated from them. Therefore, the observed variations are not caused by flux contamination from nearby sources. In addition, we argue in this paper (see Appendix D for details) that those observed variations are largely intrinsic, rather than extrinsic that is caused by refractive interstellar scintillation (RISS).

We have shown that most of the sources known to be flat-spectrum in our sample have very high brightness temperatures 10^{11-13} K ; in 4 of the objects T_{B} even exceeds the inverse Compton limit of 10^{12} K . Such high brightness temperatures indicate that the radio emission is highly Doppler beamed and hence must originate from (at least mildly) relativistic jets, as in blazars. Among the 4 with $T_{\text{B}} > 10^{12} \text{ K}$, J0849+5108 is a known OVV/HPQ blazar (e.g. Arp et al. 1979; Sitko et al. 1984, and see Appendix B) and the high brightness temperature is well expected; in turn, this gives us confidence in the results for the other objects. J0948+0022 has been previously suggested to be so by Zhou et al. (2003). For the remaining two, J1505+0326 and J1548+3511, the evidence for relativistic beaming is presented for the first time. For J1505+0326, this result is consistent with its VLBI images made at 2.3 GHz, which are dominated by a very compact component, marginally resolved at a few tens milli-arcsec resolution (Dallacasa et al. 1998). Besides, two objects, J1633+4718, and J1644+2619 have also been suggested to be blazar-like based on VLBI observations (Doi et al. 2007). Thus, in a total of 6 objects of our sample there is evidence for relativistic beaming. Assuming that the brightness temperatures of 10^{11-12} K are probably also an indication of beaming, the number of such objects will increase to 9, i.e. all the 8 flat-spectrum sources and one with unknown radio index.

Our suggestion for the presence of (at least mildly) relativistic jets is also supported by polarization detected in a few individual objects, e.g. J0849+5108 (see Appendix B) and 2MASX J0324+3410 (Neumann et al. 1994). Furthermore, this is consistent with the observed large amplitude variability in the optical in a few objects monitored, e.g. J0849+5108, J0948+0022 (Zhou et al. 2003), since normal NLS1 galaxies are found to be much less variable in the optical band (Ai et al. 2008).

These results are most naturally explained in terms of beamed emission; and thus we postulate that relativistic jets are likely present in most, if not all, of the flat-spectrum, very radio-loud NLS1 AGN. However, it should be noted that part of the observed properties above may also be explained by some other non-blazar, compact sources. For instance, flat and inverted spectra are also typical of Gigahertz-Peaked Spectrum (GPS) sources, though they generally lack of large amplitude flux variations (e.g. O’Dea 1998). Moreover, flat

spectra are often seen also in the radio sources of some ‘normal’ Seyfert galaxies, though mostly with much less power compared to the objects studied here (e.g. Ho & Ulvestad 2001). Incorporating the properties of these flat-spectrum NLS1 AGN in other wave-bands, we regard the blazar-like scenario as the most likely origin of their multi-waveband (non-thermal) emission in this paper, though other possibilities cannot be ruled out completely at this stage. We note that the majority of the sources of Komossa et al. (2006b) sample ($R_{1.4} > 10$) are steep-spectrum sources, in which strong beaming is generally not expected. The difference is largely attributed to the much higher radio-loudness cutoff of our sample, as mentioned above.

6.2.2. Optical continuum: dust extinction and jet contribution?

The majority of our sample objects (16/23) have optical continuum slopes bluer than $\alpha_\lambda = -1$. If the continuum is intrinsically blue in all objects (assuming $\alpha_\lambda = -1.7$) but is reddened by extinction, a slope of $\alpha_\lambda = -1$ corresponds to an amount of extinction $E(B-V) \simeq 0.2$, i.e. a correction factor of only ~ 2 in the B -band flux. So, even if this was the case, most of our objects would still have $R_{1.4} \gtrsim 100$, above the radio-loudness cutoff of our sample. There are 5 objects at low redshifts ($z < 0.38$) having both the $H\alpha$ and $H\beta$ lines measurable, including J1138+3653, the only one showing a significant deviation from a power-law optical continuum. J1138+3653 has a broad line Balmer decrement of $H\alpha/H\beta = 4.43$, indicative of mild extinction. Adopting the intrinsic $H\alpha/H\beta$ ratio of 3.37 [with a dispersion of 0.06 dex in $\log(H\alpha/H\beta)$] for blue radio-loud quasars (free from dust extinction) found in our recent work (Dong et al. 2008) and the Galactic extinction curve, we find an internal extinction correction $A_B \simeq 0.97$ [$E(B-V) = 0.236$], and an extinction corrected $R_{1.4}$ a factor of 2.4 lower, leading to an intrinsic radio-loudness $R_{1.4} \simeq 86$. For the remaining 4 objects, the broad line $H\alpha/H\beta$ ratios range from 2.8–3.8 with a mean of 3.4, that are well consistent with the intrinsic $H\alpha/H\beta$ distribution and hence with little or no extinction; therefore their radio-loudness $R_{1.4}$ measurements as above are almost un-affected. Particularly, J1031+4234 shows a red optical spectrum with $\alpha_\lambda = 0.55$, indicating that in at least some of the objects the red optical spectrum is intrinsic, rather than resulting from dust extinction. Another example is the blazar J0849+5108 with the reddest slope, $\alpha_\lambda = -0.20$, that is close to the previous measurement at its high light state. Its slope steepens with decreasing luminosity and changes rapidly on timescales on several months, and hence the red slope is preferentially explained as being intrinsic²³ (Arp et al. 1979), similar to red BL Lac objects (Stein et al. 1976).

Observations have revealed that in a large fraction of FSRQs the optical light can be contributed significantly by the non-thermal emission from relativistic jets that is highly beamed (e.g. Whiting et al. 2001). We have discovered that the observed optical fluxes are systematically in excess of those predicted from the broad $H\beta$ luminosity, presumably from an accretion disk, assuming that radio-loud NLS1 AGN have the same conditions in the broad line region as normal NLS1 AGN (see § 3.2). In addition, the optical slopes of our radio-loud objects are systematically bluer than that of

radio-quiet NLS1 AGN. These facts point to the presence of a second component superposed on the normal AGN-like (thermal) continuum. In light of the similar radio properties of our RL NLS1 galaxies to FSRQs, we attribute the excess optical emission to arising from the non-thermal radiation from jets, as in FSRQs, i.e. the extension to higher energies of the radio-to-infrared synchrotron radiation. This model explains naturally the observed difference in the observed-to-predicted luminosity ratios between flat- and steep-spectrum NLS1 AGN. In flat-spectrum sources, the relativistic beaming effect is strong—as suggested above—and hence the jet component over-shines the thermal optical component; while in steep-spectrum sources, the former becomes comparable to or weaker than the latter.

The observed optical continua have a variety of slopes, ranging from moderately red to blue. They are reminiscent of the optical emission of BL Lac objects, which have both red and blue continua (e.g. Stein et al. 1976), corresponding to the low- and high-peak frequency of the synchrotron hump of LBLs and HBLs, respectively. In fact, this comparison is particularly relevant in light of our finding that the RL NLS1 AGN have both LBL- and HBL-like SEDs: the blue optical continuum may be explained as contribution from the synchrotron emission peaked at high energies (UV or higher); while the red slope may be contributed by that peaked in the infrared. In turn, this is consistent with our above inference that the red slopes in our objects are mostly intrinsic, rather than resulting from extinction.

6.2.3. X-ray radiation and broad band SED

So far, we have shown that relativistically beamed jet emission, as found in blazars, provides a good explanation of the observed radio and optical properties in flat-spectrum RL NLS1 AGN. This scenario is also consistent with the fact that some of the objects show SEDs similar to FSRQs/SSRQs, while the others similar to HFSRQs in the α_{ro} , α_{ox} , and α_{rx} parameter space (Figure 9). In fact, a few of these objects were simply classified as FSRQs or SSRQs in previous studies. It is thus natural to speculate that the similar radiation mechanisms are operating in these objects as in normal FSRQs/SSRQs and HFSRQs for producing the broad-band non-thermal emission.

From the soft X-ray spectral shapes alone it is not possible to distinguish the origin of the X-ray emission, i.e. that associated with accretion disk as in normal RQ NLS1 galaxies or non-thermal jet emission as in blazars, since RQ NLS1 galaxies also show a similar Γ distribution with a large scatter (e.g. Boller et al. 1996; Grupe et al. 1999; Leighly 2000; Zhou et al. 2006). The only exception is perhaps the flat hard X-ray component ($\Gamma = 1.37 \pm 0.49$) in J1633+4718, whose best estimate value is hardly found in RQ NLS1 galaxies but is typical of FSRQs.

One way is to search for possible enhancement in the X-ray emission of RL NLS1 AGN compared to RQ objects in a statistical sense, which, if is the case, may be considered as an indirect evidence for the presence of a non-thermal jet X-ray component. Here, we simply use the X-ray detection rate in the RASS as an indicator of possible enhancement in the X-ray emission of our objects over normal NLS1 galaxies by comparing the RL and RQ samples. This is plausible since our RL NLS1 sample is essentially an optically selected one (see discussion in § 6.1), as the SDSS RQ NLS1 samples. The RASS detection rate for RQ NLS1 galaxies is $\approx 31\%$ from the SDSS DR3 NLS1 sample (Zhou et al. 2006); while it is $50(\pm 11)\%$ (11/22) for the present RL sample (excluding one

²³ Alternatively, the red slope is explained as contamination from a foreground red galaxy on the line of sight (Stickel et al. 1989), though we consider this unlikely.

solely RASS-selected target). If we consider the objects (both detections and upper limits) falling within the HBL locus only (Figure 9; excluding 2MASX J0324+3410), the detection rate is even higher $58(\pm 14)\%$ (7/12). Therefore, the higher RASS detection rate in our RL sample compared to the RQ NLS1 sample is suggestive of the presence of a significant excess X-ray component, presumably from the relativistic jets.

In Appendix E, we discuss in detail the possible mechanisms of the X-ray radiation of our RL NLS1 AGN, which are summarised here. In all of the objects, the X-rays produced in the accretion process as in normal RQ NLS1 galaxies are likely not negligible and make a considerable contribution. In addition, we suggest that the non-thermal X-ray emission from relativistic jets may also play a similarly important role. Such a component is most likely to originate from inverse Compton radiation and has a flat X-ray spectrum (in J1633+4718 and possibly in J0849+5108), as seen in FSRQs, for objects having the LFSRQ-like SEDs (NLS1 LFSRQs); whereas it is possibly the high-energy tail of synchrotron radiation with a steep spectrum for objects with the HFSRQ-like SEDs (NLS1 HFSRQs).

6.2.4. Dichotomy of LFSRQ- and HFSRQ-like NLS1 AGN

In Figure 9, there are about 7–12 objects, or 1/3–1/2 of the sample, falling within the box-shaped locus and thus having HBL/HFSRQ-like SEDs (the uncertainty comes from those with lower α_{ox} limits). Considering that radio/optically-selected (‘broad’-line) FSRQs have predominantly LBL/LFSRQ-like SED, such a high fraction of HFSRQ-like objects is surprising for our sample, which is essentially optically/radio-selected. The figure also shows that there are no objects resembling extreme HBL/HFSRQs (the lower-left corner of the HBL locus), which have the synchrotron component peaked in the X-ray regime. In fact, nearly all the candidate HFSRQ-type SEDs have the synchrotron humps peaked likely around the UV band.

Strong emission lined-FSRQs with HBL-type SEDs have recently been reported as a result of the identification of radio-X-ray (ROSAT) selected sources (Padovani et al. 1997b; Perlman et al. 1998; Padovani et al. 2003). One major distinction with our objects is that those objects have much broader lines with $FWHM > 2000 \text{ km s}^{-1}$, typical of classical FSRQs. The fraction of HBLs/HFSRQs is sample-selection dependent; the authors proposed a fraction (HBLs + HFSRQs) of about 10–15% among the whole blazar population and about 30% in the RASS-Green Bank (RGB) sample (Padovani et al. 2003). The RGB sample is comparable in the way of selection to our RASS-detected sub-sample here, which comprises 12 objects; among them, 8 objects are possibly of the HFSRQ-type (Figure 9), making the fraction as high as $\sim 2/3$. It seems that the HFSRQ fraction of our radio-loud NLS1 sample is higher than that of HFSRQs in Padovani et al. (2003).

Within the unified blazar sequence, the energy of the synchrotron peak was found to be well anti-correlated with 5 GHz radio luminosity²⁴, and with α_{rx} , which was suggested to be a tracer of the peak energy (Fossati et al. 1998). We plot in Figure 15 α_{rx} versus 5 GHz radio luminosity for the sample, with those objects without X-ray detection indicated by arrows. It clearly reveals a good correlation; the Spearman correlation test incorporating lower limits of α_{rx} using ASURV (Isobe et al. 1986; Isobe & Feigelson 1990) yields a signif-




FIG. 15.— Relationship of α_{rx} and 5 GHz radio luminosity. Symbols: arrows indicate α_{rx} upper limits for objects without X-ray detection; filled dots: flat spectrum radio sources; open circles: steep spectrum radio sources; asterisks: sources without available radio spectral indices. 2MASX J0324+3410 is also plotted (square). The dotted line corresponds to the simple division between LBLs and HBLs using α_{rx} .

icant correlation at the probability level $P_{\text{chance}}^{\text{sp}} = 0.4\%$. It can be seen that the transition between LFSRQs and HFSRQs (using $\alpha_{\text{rx}} \approx 0.78$) happens at around $\nu L_{\nu} \sim 10^{42} \text{ erg s}^{-1}$, consistent with what was previously found between HBLs and LBLs (Fossati et al. 1998, see their Figure 7). If α_{rx} does trace the synchrotron peak energy, this result shows that our radio-loud NLS1 AGN do follow the blazar sequence well. In fact, the objects at the low luminosity end are among the least radio luminous blazars known so far, falling close to or even below the low-end of the FSRQ luminosity function according to unification schemes ($L_{5\text{GHz}} \sim 10^{31.5} \text{ erg s}^{-1} \text{ Hz}^{-1}$, Urry & Padovani 1995). However, there seems to be no or at most a weak correlation between α_{rx} and $H\beta$ luminosity ($P_{\text{chance}}^{\text{sp}} = 0.1$). Such a blazar sequence also explains naturally the high fraction of HFSRQs, as the majority of our sample have lower radio powers compared to classical FSRQs. In fact, the objects of our sample have intrinsic powers mostly in the range for FR I type radio sources (see § 6.3).

6.3. Intrinsic radio power and radio-loudness

Optically thin radio emission at low frequencies is thought to be isotropic, and thus can be used as a measure of the intrinsic power of the radio sources free from Doppler boosting. The low frequency power of radio galaxies can be separated into two distinct classes at the fiducial value $L_{178\text{MHz}} \simeq 2 \times 10^{25} \text{ W Hz}^{-1} \text{ sr}^{-1}$: the FR Is with the radio emission decreasing outwards away from the nucleus, and the FR IIs which have lobes with prominent hot spots and bright edges (Fanaroff & Riley 1974). It has been suggested that RLQs are found in FR II sources, and the distinction of the FR I and FR II types is likely determined by the accretion process of the central engine (e.g. Baum et al. 1995), as well as by the density of the environment (Gopal-Krishna & Wiita 2002). Despite a lack of adequate observations on their radio morphology, it would be informative to compare the isotropic radio power of our RL NLS1 galaxies with the FR types.

We estimate the rest frame 151 MHz luminosities, $L_{151\text{MHz}}$, for our objects, based on their radio data at low frequencies,

²⁴ $L_{5\text{GHz}}$ is thought to be a good tracer of the bolometric luminosity.

which are summarised in Appendix C. For 4 objects with 151 MHz fluxes, $L_{151\text{MHz}}$ are calculated directly by using the 151–327 MHz indices for K-correction. Among the rest, 11 objects have fluxes at 327 MHz, for which $L_{151\text{MHz}}$ are estimated by extrapolating the spectra down to 151 MHz assuming the average $\alpha_{151\text{M}}^{327\text{M}} = -0.34$ found from the above 4. Using the steepest (-0.62) and the flattest ($+0.09$) $\alpha_{151\text{M}}^{327\text{M}}$ found above would increase and decrease $L_{151\text{MHz}}$ by a factor of up to 1.4 and 1.8, respectively; and thus the estimated $L_{151\text{MHz}}$ are relatively insensitive to the choice of $\alpha_{151\text{M}}^{327\text{M}}$. For the 15 objects with either detected or estimated 151 MHz fluxes, $L_{151\text{MHz}}$ range from 4.5×10^{24} to $3.5 \times 10^{27} \text{ W Hz}^{-1}$, with a median of $7.1 \times 10^{25} \text{ W Hz}^{-1}$. Among these, three have luminosities in the FR II range (J1047+4725, J1305+5116, and J1443+4725, all detected in the 151 MHz survey); while the remaining majority have similar luminosities as FR Is or are close to the transition line. With estimated $L_{151\text{MHz}} = 9 \times 10^{24} \text{ W Hz}^{-1}$, 2MASX J0324+3410 also has a luminosity typical of a FR I galaxy. However, it is not clear whether the clear separation in both radio power and morphology for radio galaxies is applicable to RL NLS1 galaxies, considering their apparently peculiar, compact morphology.

It has been suggested that, in classical RL AGN, isotropic radio emission at low frequencies can be used as a good indicator of the power of the jets (e.g. Rawlings & Saunders 1991). If this is applicable to RL NLS1 galaxies, the above results imply that our sample has much lower jet power compared to classical RLQs, which have radio power and morphology predominantly in the FR II category (e.g. Rawlings & Saunders 1991; Baum et al. 1995; Liu et al. 2006). We will address this issue in a detailed study in the future.

The radio-loudness, by its definition, is largely an apparent rather than intrinsic parameter for the sources in which the beaming effect is important above 1 GHz in the radio but less so in the optical band. As a consequence, some radio-loud AGN would have been classified as radio-intermediate or even radio-quiet, which might also possess weak and mildly relativistic jets in at least some (e.g. Falcke et al. 2006b; Blundell et al. 2003; Brunthaler et al. 2005; Wang et al. 2006), were their jets not directed close to our line-of-sight (Falcke et al. 2006a; Jarvis & Mclure 2002). A more physically meaningful parameter is the ‘intrinsic’ radio-loudness free from the beaming effect. We calculate the intrinsic radio-loudness using the un-beamed radio component, that can be extrapolated from a low-frequency flux and dominates fluxes at several GHz after the jet emission is de-boosted. The intrinsic radio-loudness at 5 GHz, $R_{5\text{GHz}}^{\text{intr}}$, is estimated by using 327 MHz fluxes for those having this flux measurement. We find $R_{5\text{GHz}}^{\text{intr}}$ in the range 34–2600 (with a median of 110) assuming a spectral slope of -0.7 for the un-beamed component, or 17–1380 (with a median of 56) assuming a slope of -1 . It should be noted that such estimates are subject to large uncertainties, partly due to the above finding that the optical light is also partially beamed in some of the objects. Nevertheless, they seem to indicate that a sizable fraction of the sample, including 2MASX J0324+3410, are likely radio-intermediate (e.g. <250 was suggested by Falcke et al. 2006a, for flat-spectrum sources). Among the sample, there are almost no genuine intrinsically RQ objects with $R \ll 10$, however.

6.4. Black hole masses of radio-loud NLS1 galaxies

6.4.1. Estimation of black hole mass and Eddington ratio

It has become possible recently to estimate the black hole masses M_{BH} of AGN from single epoch spectroscopic observational data (e.g. Kaspi et al. 2000; Wandel et al. 1999), though the uncertainties of both systematics and scatter remain large (Vestergaard & Peterson 2006; Collin et al. 2006). Assuming that the motion of the line-emitting clouds are virialised, the black hole mass can be determined by $M_{\text{BH}} = f R_{\text{BLR}} \Delta v^2 / G$, where f is a scale factor depending on the kinematics and geometry of the BLR, Δv the emission line width, and R_{BLR} the BLR radius. We assume that the commonly used R_{BLR} –luminosity relation, derived via reverberation mapping for nearby, normal AGN, is applicable to NLS1 AGN. We use the relation for the $\text{H}\beta$ luminosity given by Kaspi et al. (2005, their Eq. 1 and Table 3); we intend not to use the continuum 5100 Å luminosity for possible contamination from jet emission (see Section 3.2). A common practice, among a few others²⁵, is to take $FWHM$ as Δv and $f = (\sqrt{3}/2)^2 = 0.75$, assuming an isotropic distribution with random orbital inclinations of the broad line clouds (Netzer 1990). We follow the practice in this work, the same as in Komossa et al. (2006b).

The estimated black hole masses of our sample range from 1×10^6 to $1.5 \times 10^8 M_{\odot}$, peaked at near the median $1.3 \times 10^7 M_{\odot}$. The black hole mass distribution is well consistent with that of the Komossa et al. (2006b) sample. However, by studying the systematic effects in M_{BH} estimation, Collin et al. (2006) argued that the use of $FWHM$ rather than line-dispersion (σ_{line}) would introduce systematic bias in the scaling factor f . We also give alternative M_{BH} estimation by correcting this effect using the formalism proposed by Collin et al. (2006, their eq.7). For the objects in our sample ($FWHM < 2200 \text{ km s}^{-1}$) specifically, this correction sets a value of the scale factor $f \simeq 1.5$, about twice the value in the former estimation. This results in M_{BH} estimates a factor of 2 larger than the values obtained above, ranging from 2×10^6 to $3 \times 10^8 M_{\odot}$ with a median $2.6 \times 10^7 M_{\odot}$. The black hole masses thus estimated are listed in Table 4, which are used in the following analysis.

Having estimated the masses of the central black holes, now we can derive the Eddington ratios, R_{Edd} . For bolometric luminosity correction, we assume $L_{\text{bol}} = 9 \lambda L_{\lambda 5100}$, as suggested by, e.g. Warner et al. (2004), Vestergaard (2004), and Kaspi et al. (2000). The $\lambda L_{\lambda 5100}$ luminosity is estimated from the broad $\text{H}\beta$ line luminosity using Eq.(5) of (Zhou et al. 2006) to eliminate the effect of the extra jet contribution. It should be noted that such an estimate is subject to large systematic uncertainties inherent in estimation of M_{BH} , as well as of bolometric luminosity. Using the M_{BH} values obtained with the scale factor $f = 0.75$ (see above), we find the Eddington ratios ranging from $R_{\text{Edd}} = 0.52$ to 2.6 and peaked around a mean of 1.0, similar to those found by Komossa et al. (2006b) with a mean of 1.2. However, when the correction of Collin et al. (2006) is applied, the M_{BH} values become about twice larger and, consequently, the obtained R_{Edd} values become about twice smaller, i.e. ranging from 0.27 to 1.28 with an average of 0.52.

Now we compare the distributions of M_{BH} and R_{Edd} between our RL sample and the normal NLS1 sample of Zhou’06. Our RL sample has significantly higher M_{BH} and

²⁵ A few other scaling relations for M_{BH} estimation were recently presented in the literature (e.g. Onken et al. 2004; Greene & Ho 2005, 2007; Vestergaard & Peterson 2006; Collin et al. 2006), that slightly differ from one another.

R_{Edd} distributions compared to the RQ one. This agrees with the general trend that, as aforementioned in § 6.1, RL objects are more likely to be found in AGN with massive black holes (e.g. Best et al. 2005). Given their narrow ranges of the line-width, the black hole masses of NLS1 samples, as determined from the $H\beta/H\alpha$ line-width and luminosity, is strongly correlated with luminosity, and in turn, with redshift for flux limited samples. This may explain the higher redshift distribution of our RL sample compared to that of the normal NLS1 sample of Zhou'06 (see Figure 2).

It should be pointed out that the virial M_{BH} estimation is subject to assumption of an isotropic geometry of the BLR with random orbital inclination of the clouds. If the BLR is flat, a small line width would be a result of small inclination, and then M_{BH} would be largely underestimated (see, e.g. Wills & Browne 1986; Decarli et al. 2008). Indeed there are some suggestions in the literature for a flattened geometry of the BLR in some RL AGN (e.g. Vestergaard et al. 2000; Sulentic et al. 2003; Jarvis & McLure 2006); but see Punsly (2007). This caveat concerns particularly those of our objects, that have compact radio morphology and those that are blazar-like, indicating a small viewing angle. Given that the M_{BH} distribution of our sample is indistinguishable from that of the RQ NLS1 sample (within the same redshift range), this comes back to the old question whether NLS1 galaxies, as a population, have flattened BLRs and are viewed preferentially face-on, rather than having small M_{BH} (e.g. Osterbrock & Pogge 1985; Puchnarewicz et al. 1992; Bian & Zhao 2004); but see e.g. Smith et al. (2004).

If RL NLS1 galaxies do possess relativistic jets and hence anisotropic radio emission, as we have shown here, they can be used as a probe to this question since their inclination can be inferred from radio observations in a statistic manner. However, with the relatively few data available currently, we do not find any statistically significant trends of the dependence of line-width (M_{BH}) on probable inclination indicators, such as radio-loudness or radio slope. Therefore, though there is no compelling evidence supporting the planar BLR nature of RL NLS1 from the present data, we cannot rule out this possibility, either. A much larger RL NLS1 sample covering a wide range of radio-loudness is desirable to shed light to this important question in the future.

6.4.2. Radio-loudness versus black hole mass

As one of the fundamental relations in understanding AGN jets, the dependence of radio-loudness R on M_{BH} has been studied extensively in recent years, with the advent of the easy estimation of M_{BH} in AGN. A R - M_{BH} correlation was proposed by Laor (2000) that all RLQs in the PG quasar sample have $M_{\text{BH}} > 3 \times 10^8 M_{\odot}$, while the radio-quiet quasars have a much lower M_{BH} distribution. This trend was confirmed in some studies (e.g. Lacy 2001; McLure & Jarvis 2004; Metcalf & Magliocchetti 2006), but was found to be weak or absent in others (Oshlack et al. 2002; Woo & Urry 2002). Furthermore, by including Seyferts and low luminosity AGN, Ho (2002) found that R is mostly correlated with R_{Edd} rather than M_{BH} . In a recent comprehensive treatment of this question, Sikora et al. (2007) demonstrated that R depends on both R_{Edd} and M_{BH} , with two distinct inverse R - R_{Edd} dependence sequences for AGN hosted by elliptical (systematically higher R) and disk galaxies (systematically lower R), respectively. Regardless whether there is indeed a trend of dependence or not, most aforementioned studies show that, on the M_{BH} - R plane, there is a scarcity of RL AGN with low M_{BH} . Even for those low-




FIG. 16.— Dependence of radio-loudness on black hole mass for the radio-loud NLS1 galaxies of our sample (filled dots) and those from the sample of Komossa et al. (2006b, open squares). Over-plotted are schematic areas populated by the bulk of the radio sources of the samples of Laor (2000, solid line) and Lacy (2001, dashed line). We note that a small number of (normal) RL AGN with small M_{BH} and large R has also been claimed by Woo & Urry (2002), which lies mostly at higher $R \simeq 10^{3-4}$ above ours in the diagram; however, there remains a controversy as to the estimation of those M_{BH} (see, e.g. Jarvis & McLure 2002; Laor 2003).

M_{BH} RL AGN suggested in the literature (e.g. Oshlack et al. 2002; Woo & Urry 2002), there remains a controversy as to the estimation of M_{BH} and R (see, e.g. Jarvis & McLure 2002; Laor 2003).

Obviously, as first pointed out by Wang et al. (2001) and Zhou et al. (2003), radio-loud NLS1 AGN do *not* follow the above trend found for normal RL AGN (provided the virial M_{BH} as estimated above are largely correct). Moreover, Komossa et al. (2006b) demonstrated explicitly that RL NLS1 galaxies are located in a region of the M_{BH} - R diagram that is sparsely populated by other types of AGN ($R \simeq 10^{-1-3}$ and $M_{\text{BH}} < 10^8 M_{\odot}$).

Similarly, we plot $R_{1.4}$ versus M_{BH} for our very RL NLS1 galaxies in Figure 16. Also plotted are the loci of the objects in the samples of Laor (2000) and Lacy (2001) for comparison. Two pieces of results can be inferred immediately from Figure 16. Firstly, within our sample, there is no correlation found between $R_{1.4}$ and M_{BH} , nor between $R_{1.4}$ and R_{Edd} ($P_{\text{chance}}^{\text{sp}} = 0.44$ and 0.27 , respectively); however, we note that these results are rather uncertain given the small sample size and the limited ranges in both $R_{1.4}$ and R_{Edd} . Secondly, our new result confirms and strengthens that of Komossa et al. (2006b), and improves upon it by extending the RL NLS1 locus to even lower M_{BH} and higher $R_{1.4}$. In particular, the bulk of our RL NLS1 galaxies lies within a locus of $M_{\text{BH}} \simeq 10^{6-7.5} M_{\odot}$ and $R \simeq 10^{-1-3}$, which appears to be preferentially avoided by other types of RL AGN [compared to, e.g. Sikora et al. (2007), see their Figure 4], except for low power Seyfert galaxies and LINERs; however, those exceptions have exclusively low Eddington ratios $R_{\text{Edd}} < 10^{-2}$ and are drastically different from NLS1 with high R_{Edd} (see also Ho 2002). Compared to Figure 4 in Sikora et al. (2007), our RL NLS1 galaxies appear to have the largest radio-loudness up to $R_{1.4} \sim 10^{3-4}$ among all types of AGN within this M_{BH} range. This means that the RL NLS1 galaxies show strong excess radio emission with respect to their M_{BH} , when the suggested radio power- M_{BH} relation for normal AGN is compared²⁶ (e.g. Lacy 2001). Finally, we note that the above

²⁶ We note that several (normal) RL AGN with small M_{BH} similar to ours and with even larger R ($\simeq 10^{3-4}$) have also been claimed by Woo & Urry (2002), though there remains a controversy as to the estimation of M_{BH} (see,

two results remain valid even if the Doppler de-boosted radio-loudness values (see § 6.3) are used instead, which are in the range of $R \simeq 10^{1-3}$.

Hence, the existence of RL NLS1 galaxies as a *population* with small M_{BH} , if estimated correctly, fills up the previously sparsely populated region in the $M_{\text{BH}}-R$ parameter space for normal RL AGN. This result indicates that there indeed exist AGN with high radio-loudness ($R \sim 10^{2-3}$) but intermediate M_{BH} ($10^{6-8} M_{\odot}$). Though there is the systematic uncertainty in estimating M_{BH} ascribed to the possible BLR inclination effect, we argue that this result may not be affected qualitatively. This is because the Komossa et al. sample, though composed of mainly steep spectrum sources (thus relatively larger inclination), led to similar results. Furthermore, in a few objects where the host galaxies are resolved, their small M_{BH} are supported by a lack of massive galactic bulges, as predicted by the M_{BH} –bulge relation (e.g. Magorrian et al. 1998). Our result also implies that the inverse relation between R and R_{Edd} as suggested previously (see, e.g. Greene et al. 2006; Sikora et al. 2007) may not work at the high R_{Edd} end close to the Eddington rate. Furthermore, Sikora et al. (2007) argued that RL AGN hosted by elliptical galaxies are much radio-louder than AGN hosted by disk galaxies. Though this statement may be true in general, it is perhaps more appropriate in terms of RL fraction, i.e. the fraction of RL objects is lower in AGN hosted by disk galaxies compared to that of AGN hosted by ellipticals; in addition, the difference in radio-loudness is now less than what was thought previously (Sikora et al. 2007).

6.5. On the nature of radio-loud NLS1 galaxies

6.5.1. A previously unknown population of radio-loud AGN?

On the black hole mass–radio-loudness plane our RL NLS1 galaxies form a natural extension to low M_{BH} of classical RLQ/BLRGs (Broad Line Radio Galaxies) with roughly constant radio-loudness (Figure 16; see also Figure 3 in Sikora et al. 2007). This is not surprising, since the transition between BLAGN and NLS1 AGN is *smooth* in the line-width of the broad lines, and, as a consequence, in black hole mass and luminosity. If the M_{BH} are underestimated systematically (see § 6.4), the loci of the radio-loud NLS1 AGN will move even closer to those of normal RL AGN. In this regard, it is natural to treat RL NLS1 galaxies as the low- M_{BH} end of normal RL AGN.

On the other hand, however, RL NLS1 galaxies show some peculiarities in observed properties that are unusual in comparison with, or at an extreme of, normal RL AGN. The most distinctive features are in the radio properties. Even for steep-spectrum sources, the radio sources of RL NLS1 galaxies appear to be compact, with the size less than several tens of kpc or even smaller wherever higher resolution observations are available. In contrast, the radio sources of RLQ/BLRGs are about 100–400 kpc in a similar redshift and luminosity range as our radio-loud NLS1 AGN (Singal 1993). Moreover, most of the RL NLS1 galaxies have radio power in the range of FR I radio galaxies, whereas RLQ/BLRGs tend to be associated with FR II type sources²⁷ (e.g. Laing et al. 1983).

In the following aspects the two types are not mutually exclusive but overlap to some extent, yet they seem to differ in a systematic way. Firstly, There seems to be a higher fraction

of HBL/HFSRQ-like blazars in optically selected RL NLS1 galaxies than in normal RLQ/BLRGs of radio/optically selected samples (Padovani et al. 2003), though further confirmation using complete samples is needed. Secondly, as commonly suggested, NLS1 AGN are accreting at close to, or larger than the Eddington limit (e.g. Collin & Kawaguchi 2004). At such high rates the accretion flow proceeds via a slim disk where energy is partially advected into the black hole before being radiated due to relatively long cooling time (e.g. Abramowicz et al. 1988; Mineshige et al. 2000; Wang & Netzer 2003). Thus the accretion processes in radio-loud NLS1 AGN may differ drastically from those in normal RLQ/BLRGs, where a standard thin disk is believed to be operating, though they do overlap at $R_{\text{Edd}} \simeq 1$ to some extent (see e.g. Sikora et al. 2007). The final point—on the host galaxies—is only suggestive because of a lack of sufficient evidence. RL NLS1 nuclei, at least some, seem to be hosted in disk or interacting galaxies, and/or galaxies with relatively young stellar populations. In contrast, classical RLQ/BLRGs²⁸ are preferentially hosted in giant elliptical/massive bulge dominated galaxies or at least strongly passively evolving galaxies (e.g. Bahcall et al. 1997; Dunlop et al. 2003; Floyd et al. 2004; O’Dowd & Urry 2005), although there was also evidence reported for star-formation in some hosts beyond $z \gtrsim 0.4$ (e.g. Örm Dahl & Rönnback 2005; O’Dowd & Urry 2005).

In consideration of their unusual properties, we are left with two options regarding the nature of RL NLS1 galaxies: either they are a downsized extension of normal RL AGN to the lower- M_{BH} (and perhaps higher- R_{Edd}) range, or they form a previously unknown population distinct from normal RL AGN. The former would invoke some unknown processes that can explain the observed differences between RL NLS1 and normal RL AGN with the smooth change of M_{BH} . For the latter, the operating physical processes in likely black hole accretion and jet formation may be radically different for the two classes. A related, interesting question is whether the differences in the observed properties between RL NLS1 and normal RL AGN can be naturally explained by the same underlying physical processes that make NLS1 different from normal AGN, or whether some additional mechanisms are needed. To further understand these questions, a large sample of RL NLS1 galaxies with the full range of radio-loudness is needed, as well as extensive observations, especially in the radio band. As complementary discussion, we also compare them with two special types of RL AGN below, namely, BL Lac objects and compact steep-spectrum sources. As will be seen, RL NLS1 galaxies may be a heterogeneous population themselves.

6.5.2. Relationship with BL Lac objects

One object in our sample, J0849+5108 (0846+51W1), was formerly classified as a BL Lac object at its high light state based on previous observations (see Appendix B for details). As such, it presents an interesting case linking possibly RL NLS1 galaxies with BL Lac objects, as pointed out by Zhou et al. (2005), though its black hole mass ($2.6 \times 10^7 M_{\odot}$) is lower compared to those typical of BL Lac objects ($M_{\text{BH}} \sim 10^{8-9} M_{\odot}$, e.g. Cao 2003). In fact, it has been speculated in

e.g. Laor 2003). Our sample appears to differ from theirs in the sense that it fills up the gap between $R \simeq 10$ and 10^3 in the small M_{BH} regime in their $M_{\text{BH}}-R$ diagram.

²⁷ There are exceptions, though; see, e.g. Blundell & Rawlings (2001).

²⁸ Ho & Peng (2001) showed that some Seyfert galaxies hosted in disk-dominated spirals are in fact RL when the optical nuclear luminosities are considered properly. However, the vast majority of their sample objects are of substantially low power and sub-Eddington (Ho 2002).

the past decade that concealed Seyfert nuclei exist in a few BL Lac objects, including BL Lac itself (Vermeulen et al. 1995; Corbett et al. 2000). It is therefore tempting to postulate that some RL NLS1 galaxies may actually be BL Lac objects (with low M_{BH}) at their low light states.

The bulk of our RL NLS1 galaxies have a BLR luminosity²⁹ distribution lying within $\log(L_{\text{BLR}}/\text{erg s}^{-1}) = 43.5\text{--}44.5$, the low end of which overlaps partially with those of BL Lac objects (a median of $10^{43.7} \text{ erg s}^{-1}$ was given for the upper limits of L_{BLR} by Celotti et al. 1997). Specifically, J0849+5108 has $L_{\text{BLR}} \sim 10^{43.7} \text{ erg s}^{-1}$. However, this is likely not the case for most of the RL NLS1 galaxies given their relatively higher L_{BLR} compared to the upper limits for BL Lac objects. On the other hand, this possibility means that BL Lac objects might not be a homogeneous population, and some may just be Seyfert nuclei/NLS1 at a high state of the non-thermal radiation from the jet. Similarly, this concerns only BL Lac objects at the low- M_{BH} end, as is the case of J0849+5108, rather than the bulk of classical BL Lac objects that harbour massive black holes and accrete likely via a low efficiency accretion flow (e.g. Cao 2003).

6.5.3. Relationship with CSS sources

Compact steep-spectrum sources are compact, powerful radio sources with spectral peaks near 100 MHz. They are inherently compact with linear sizes ≤ 20 kpc, having (often symmetric) double/triple morphology with complete jets and lobes on small scales. The most popular scenarios to explain their nature advocate that they are either young and evolving radio sources or confined by dense gas of their environment (see O’Dea 1998, for a review). Very recently, a few lines of observational evidence point to an interesting link between RL NLS1 galaxies and CSS, which share some degree of similarity as the following. (1) As mentioned in § 1, the radio sources of all RL NLS1 galaxies are compact, including steep-spectrum sources. In particular, a few NLS1 AGN having higher resolution observations show small-size structure: e.g. most of the radio emission of 2MASX J0324+3410 is from a region smaller than 6.8 mas (8.2 pc projected size), as seen from VLBA images at 8.4 GHz (in preparation). (2) The low frequency spectrum of J1047+4725, one of the steep-spectrum objects in our sample, appears to flatten at around a few hundred MHz as indicated by its flat slope of -0.28 between 327 and 151 MHz; such a spectral break is characteristic of CSS. In fact, the RL NLS1–CSS connection was advocated by Oshlack et al. (2001) and Gallo et al. (2006) for the case of PKS 2004-447, although it is unclear as to whether PKS 2004-447 is a genuine NLS1 galaxy (see Zhou et al. 2003, for a short comment). (3) There are observational indications that the host galaxies of a large fraction of CSS are interacting/merging and may contain significant amount of dense gas (O’Dea 1998), similar to what was found in some RL NLS1 galaxies.

The radio power distribution of our RL NLS1 galaxies overlaps with the lower part of that of CSS (O’Dea 1998). As such, if there is an overlap between CSS and RL NLS1 galaxies, they should be mostly the CSS having low radio power, and, relatively small M_{BH} . Since CSS sources also possess (small scale) jets, in cases where the jets are directed close to

our line-of-sight, we expect to see blazar-like objects. Some of those could be the flat-spectrum RL NLS1 galaxies studied here, in which relativistic jets are inferred to be present. Conclusive remarks have to await future proper and detailed comparisons of the two classes, as well as more radio observations with higher resolution, however.

6.5.4. Implication for jet formation in AGN

Given the high Eddington ratios of NLS1 galaxies in general, as commonly suggested and also found here, our RL NLS1 AGN present a strong case where jets, possibly at least mildly relativistic, can be produced in high R_{Edd} systems. This strengthens the similar situation in powerful RLQs with much larger M_{BH} . This contrasts clearly with black hole X-ray binaries where jets are found at the low/hard state, despite the suggestion that transient jets (or mass ejection) may occur at the so called very-high state where R_{Edd} is higher (Maccarone et al. 2003; Fender 2004). The formation of jets is thought to be related to the black hole and accretion process (e.g. Blandford 2000; Celott & Blandford 2001; Meier 2003; Ballantyne 2007). The finding of relativistic jets in black hole accretion systems with high R_{Edd} (where the accretion proceeds via probably a new form other than the standard disk) and, moreover, with relatively low M_{BH} is important for addressing basic questions as to how jets are formed. Jet formation in high R_{Edd} systems (slim disks) is far from clear, though there are some suggestions that outflows as well as mildly beamed and collimated emission are likely present in such systems (Ohsuga et al. 2005).

As has been shown above (see also Komossa et al. 2006b; Whalen et al. 2006), apart from the continuum radiation, the RL NLS1 galaxies resemble closely their RQ counterparts in many ways, e.g. optical emission properties (except for stronger Fe II emission), the distributions of black hole masses and the Eddington ratios. These facts suggest that RL NLS1 galaxies are simply otherwise normal NLS1 galaxies but having relativistic jets, reminiscent of the RL/RQ dichotomy of more powerful, classical AGN. It would be interesting to find out whether there is a bimodal distribution in the radio-loudness, as reported for the broad-line quasar population (though this is still a matter of debate), in future work considering objects with the whole range of radio-loudness. The nearly indistinguishable properties other than the presence of radio jets possessed by the two subsets of NLS1 galaxies, RL and RQ, seemingly hint at a possibly important role of black hole spin in the making of relativistic jets in RL NLS1 galaxies.

7. SUMMARY OF CONCLUSIONS AND IMPLICATIONS

A homogeneous sample of genuine RL NLS1 galaxies (radio-loudness $R_{1.4} \gtrsim 100$) is presented for the first time, comprising 23 objects drawn from SDSS spectroscopic objects with FIRST detections. Their NLS1 nature is discovered by analysing systematically the spectroscopic data from the QSO and galaxy database in the SDSS DR5. The sample can be treated as an optically–radio selected one. We carry out systematic investigations into their multiwaveband properties from radio to X-rays using data available from the archives. The main results are the following.

All radio sources of the sample are compact, less than a few tens of kilo-parsec at least. Several with VLBI observations are unresolved or marginally resolved at scales of about tens of parsec. Among those with available radio spectral indices,

²⁹ The luminosity of the broad line region is estimated from the $H\beta$ luminosity using the method proposed by Celotti et al. (1997) and the quasar emission line ratios given by Francis et al. (1991), yielding $L_{\text{BLR}} = 25.26 L_{H\beta}$ (see also Liu et al. 2006).

the majority are flat-spectrum sources ($\alpha > -0.5$ around 1.4–5 GHz), and a few even have inverted spectra ($\alpha > 0$). Radio variability is found to be common in most of the flat-spectrum sources. We were able to estimate the brightness temperatures of the radio sources, that are as high as $T_B > 10^{11}$ K and even exceeding the inverse Compton limit (10^{12} K) in four objects. The high brightness temperatures indicate the presence of relativistic beaming. The majority of RL NLS1 galaxies has a low-frequency (isotropic) radio power falling in the range of FR I type radio galaxies ($L_{151\text{MHz}} \lesssim 10^{25} \text{ W Hz}^{-1} \text{ sr}^{-1}$), and the remaining minority in the range of the FR II type. When isotropic, rather than Doppler boosted, emission is considered, a large fraction of the objects can be considered as intrinsically radio-intermediate.

The observed optical luminosities are systematically in excess of the thermal continua predicted from the $H\beta$ line luminosities by a factor of up to several. The optical slopes of our RL objects, while spanning a wide range from moderately red to blue, are systematically bluer than that of radio-quiet NLS1 AGN. We interpret this effect as the contribution from a (non-thermal) component, presumably from relativistic jets, to the optical band superimposing the central ionising continuum seen in radio-quiet NLS1 AGN. The optical Fe II emission in the radio-loud NLS1 in our sample is on average stronger than ‘normal’ NLS1 galaxies. Except this, the two subsets of NLS1 galaxies seem to be indistinguishable in other emission line properties.

On the radio–optical–X-ray effective spectral indices plane (α_{ox} vs. α_{ro}), our objects have broad-band SEDs well consistent with those of blazars, i.e. both the HFSRQ and LFSRQ types as well as SSRQs. Of particular interest, the fraction of the candidate HFSRQ-type objects seems to be high, though their synchrotron component peaks are only moderately high (at around the UV). This finding extends the recently discovered strongly-lined (FSRQ) HBL-type blazars (Padovani et al. 2003) to NLS1 galaxies.

The continuum shape in the soft X-ray band exhibits a large variety among our RL NLS1 galaxies. There is evidence in one, or possibly two objects for a flat X-ray spectral component ($\Gamma \lesssim 1.8$), which we interpret as an inverse-Compton origin from jets as in FSRQs. Most of the objects with estimation of the X-ray spectral slopes show steep spectra. The steep X-ray spectrum is also consistent with what is expected for HFSRQs, that originates from the high energy end of the synchrotron component. Based on this circumstantial evidence, we suggest that such a component may be a significant contributor to the observed X-ray emission, though the X-ray component as seen in radio-quiet NLS1 galaxies may also play a role.

In view of the independent lines of evidence presented here, as well as those found in previous studies for several individuals (either inclusive or exclusive of our sample), we suggest that there likely exists a population of RL NLS1 galaxies showing the broad-band continuum emission characteristic of blazars, though alternative explanations cannot be ruled out completely. In this scenario, beamed non-thermal emission from jets, that are at least mildly relativistic, dominates the high-frequency radio band, and contributes significantly to the optical and possibly the X-ray bands. Our finding extends the radio-loud quasar population into a narrow line-width regime of the broad emission lines. We postulate that all genuine radio-loud NLS1 galaxies possess small scale, at least mildly relativistic jets, reminiscent of classical radio-loud AGN. In this scenario, steep-spectrum NLS1 sources (perhaps includ-

ing some of those with smaller radio-loudness within $R=10$ –100) are simply objects of the same population whose radio jets are aligned at a relatively large angle to the line-of-sight.

The host galaxies of two nearby RL NLS1 nuclei, whose images are resolved, have disk morphology, one even in a merger, in line with the finding for the previously published blazar-NLS1 hybrid object 2MASX J0324+3410 (Zhou et al. 2007). In several objects imprints of young stellar populations are likely to be present in form of high-order Balmer absorption lines in the SDSS spectra.

Based on the commonly used relation of black hole mass with broad line width and luminosity, our RL NLS1 galaxies have estimated M_{BH} in the range of 10^6 – $10^8 M_{\odot}$, provided that this M_{BH} scaling relation is applicable to NLS1 and the BLR is not largely flattened. The thus estimated Eddington ratios range from $R_{\text{Edd}}=0.5$ to 3, and peak at $R_{\text{Edd}} \approx 1$ (assuming $L_{\text{bol}} = 9\lambda L_{\lambda 5100}$). RL NLS1 galaxies deviate from the known trends of radio-loudness versus the Eddington ratio and black hole mass, and are populated in a region in the radio-loudness– M_{BH} diagram where normal AGN are seldom found, i.e. moderate M_{BH} and large R ($M_{\text{BH}} \simeq 10^{6-7.5} M_{\odot}$, $R \simeq 10^{1-3}$), confirming and improving upon the previous results suggested by Komossa et al. (2006b). As such, RL NLS1 show strong excess radio emission with respect to their M_{BH} , when compared to the previously suggested radio power– M_{BH} relation for normal AGN (e.g. Lacy 2001). Our results imply that large radio-loudness ($R = 10^{1-3}$) can be produced in AGN with moderate M_{BH} down to $\sim 10^6$ – $10^7 M_{\odot}$ and accreting at/close to the Eddington rate; and some of these AGN are hosted in late-type galaxies, some even in interacting systems. The presence of relativistic jets in such systems may have interesting implications for understanding jet formation in AGN in general.

In spite of a likeness to normal RL AGN in many aspects, RL NLS1 galaxies show some extreme or unusual properties, though some of the evidence are only circumstantial. These include compact radio morphology, relatively low radio power, a high incidence of HBL/HFSRQ-type SEDs, host galaxies of probably late-type and/or with young stellar populations, and the common properties characteristic of NLS1 galaxies, e.g. smaller black hole masses, higher Eddington ratios, and stronger Fe II emission. All of these lead us to postulate that either they are an extension of normal RL AGN to the lower- M_{BH} range, and all those differences can be explained as a consequence of decreasing M_{BH} ; or they form a previously unknown population distinct from normal radio-loud AGN. As one interesting possibility, we speculate that low-power CSS might be the parent population of the flat-spectrum RL NLS1 galaxies with beamed jet emission. For the future, a much larger sample and extensive multi-waveband observations are essential for understanding the nature of radio-loud NLS1 galaxies.

Finally, if the broad-band radiation mechanisms in flat-spectrum radio-loud NLS1 AGN are indeed the same as in blazars, we expect some of these objects to be γ -ray emitters in the GeV band. This can be tested by observations of GLAST in the near future.

We thank the anonymous referee for helpful comments and suggestions. This work is supported by the Chinese Natural Science Foundation through projects No. NSF10533050 and NSF10473013, and the BaiRen Project of the Chinese Academy of Sciences. Funding for the creation and the dis-

tribution of the SDSS Archive has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Aeronautics and Space Administration, the National Science Foundation, the U.S. Department of Energy, the Japanese Monbukagakusho, and the Max Planck Society. The SDSS is managed by the Astrophysical Research Consortium (ARC) for the Participating Institutions. The Participating Institutions are The University of Chicago, Fermilab, the Institute for Advanced Study, the Japan Participation Group, The

Johns Hopkins University, Los Alamos National Laboratory, the Max-Planck-Institute for Astronomy (MPIA), the Max-Planck-Institute for Astrophysics (MPA), New Mexico State University, Princeton University, the United States Naval Observatory, and the University of Washington. This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

APPENDIX

A. OPTICAL SPECTRAL MODELING

The starlight and nuclear continua are modeled as

$$S(\lambda) = A_{\text{host}}(E_{B-V}^{\text{host}}, \lambda) A(\lambda) + A_{\text{nucleus}}(E_{B-V}^{\text{nucleus}}, \lambda) [bB(\lambda) + c_b C_b(\lambda) + c_n C_n(\lambda) + dD(\lambda)] \quad (\text{A1})$$

where $S(\lambda)$ is the observed spectrum. $A(\lambda) = \sum_{i=1}^6 a_i IC_i(\lambda, \sigma_*)$ represents the starlight component modeled by our 6 synthesized galaxy templates, which had been built up from the spectral template library of Simple Stellar Populations (SSPs) of Bruzual & Charlot (2003) using our new method based on the Ensemble Learning Independent Component Analysis (EL-ICA) algorithm. The details of the galaxy templates and their applications are presented in (Lu et al. 2006). $A(\lambda)$ was broadened by convolving with a Gaussian of width σ_* to match the stellar velocity dispersion of the host galaxy; in this process the velocity difference between emission and absorption lines is taken into account. The un-reddened nuclear continuum is assumed to be $B(\lambda) = \lambda^{-1.7}$ as given in Francis (1996). We modeled the optical Fe II emission, both broad and narrow, using the spectral data of the Fe II multiplets for I Zw I in the $\lambda\lambda 3535\text{--}7530\text{\AA}$ range provided by Véron-Cetty et al. (2004, Table A1,A2). We assume that the broad Fe II lines (C_b in Eq.1) have the same profile as the broad $H\beta$ line, and the narrow Fe II lines (C_n), both permitted and forbidden, have the same profile as the that of the narrow $H\beta$ component, or of [O III] $\lambda 5007$ if $H\beta$ is weak. $D(\lambda)$ represents the templates of higher-order Balmer emission lines ($10 \leq n \leq 50$) and Balmer continuum generated in the same way as in Dietrich et al. (2003). $A_{\text{host}}(E_{B-V}^{\text{host}}, \lambda)$ and $A_{\text{nucleus}}(E_{B-V}^{\text{nucleus}}, \lambda)$ are the color excesses due to possible extinction of the host galaxy and the nuclear region, respectively, assuming the extinction curve for the Small Magellanic Cloud of Pei (1992). The fitting is performed by minimizing the χ^2 with E_{B-V}^{host} , E_{B-V}^{nucleus} , a_i , σ_* , b , c_b , c_n , and d being free parameters.

B. NOTES ON INDIVIDUAL OBJECTS

J0849+5108 (0846+51W1): This object has drawn substantial attention ever since its first identification with a BL Lac object (at a bright state) by Arp et al. (1979), at a claimed high redshift of $z = 1.86^{30}$. Its radio flux also showed large amplitude variations up to 50% at 1.4 GHz. This object exhibited a remarkable optical brightening with an amplitude of $\Delta V \sim 5$ within less than a month (Arp et al. 1979) and high and variable optical polarization (Moore & Stockman 1981; Sitko et al. 1984), making it one of the optically violent variables (OVV) and highly polarized QSOs (HPQ). Night-to-night and even intra-night optical variability was also found (Sitko et al. 1984; Stalin et al. 2005). At maximum light outburst, the continuum is featureless, whereas its spectra taken at normal/minimum light states show strong emission lines (Arp et al. 1979; Stickel et al. 1989). Thus it is a transition between a quasar and a BL Lac object, i.e. appeared to be a BL Lac object only at its flaring state. The large and variable polarization in the optical band suggests strongly that this emission is due to synchrotron radiation (Sitko et al. 1984). The conspicuous characteristic emission lines identified in the SDSS spectrum of J0849+5108 give a redshift of $z = 0.583$. A detailed study of its SDSS spectrum and other properties was presented by Zhou et al. (2005). The optical spectral slope obtained from fitting the SDSS spectrum is red, with $\alpha_\lambda \simeq -0.2$.

J0948+0022: The object was first discovered for its radio emission in various radio surveys ($S_{5\text{GHz}} \sim 0.2\text{--}0.3\text{Jy}$). It was identified with a NLS1 galaxy, based on its SDSS spectrum, by Williams et al. (2002) and Zhou et al. (2003), and was studied in detail as one of the few previously known very RL NLS1 galaxies. The radio source has an inverted radio spectrum and was unresolved with VLBI at milli-arcsec resolution. These observations were explained in terms of relativistic beaming (Doi et al. 2006). A very high brightness temperature ($\sim 10^{13}\text{ K}$) inferred from radio flux variations suggest the presence of a jet with relativistic beaming (Zhou et al. 2003). Long-term optical variability with an amplitude of about 1 mag in the B -band was also noted in Zhou et al. (2003).

J1505+0326: It has radio fluxes peaked at $\sim 7\text{ GHz}$ (Tinti et al. 2005). Its VLBI images are dominated by a very compact component, marginally resolved at 2.3 GHz band (Dallacasa et al. 1998). The radio emission at centimeter wavelengths shows variability.

J1633+4718 (RXJ 16333+4719): The object was identified with a NLS1 galaxy in optical identification of ROSAT All-Sky Survey (RASS) sources (Moran et al. 1996; Wisotzki & Bade 1997). The AGN (see Figure 1 for its SDSS spectrum) is actually one of the two nuclei, separated by $4''$, residing in an interacting/merger system. The other nucleus has a starburst spectrum (Bade et al. 1995). The radio source, which is associated with the AGN, is unresolved with VLBI at milli-arcsec resolution (Doi et al. 2007). Its spectral shape above 5 GHz is inverted, as measured by simultaneous multiwavelength observations by Neumann et al. (1994). The system was noticed to be associated with the IRAS source IRAS 16319+4725 (RA=16h33m23.03s, Dec=+47d19m03.5s), whose positional uncertainty—much larger than the separation of the two optical nuclei—encompasses

³⁰ This redshift value, given by the authors based on the spectra with low resolution and signal-to-noise ratios, turned out to be wrong.

both of the nuclei (Boller et al. 1992; Moran et al. 1996; Condon et al. 1998b). The [O III] doublet shows a noticeable blue wing in its SDSS spectrum.

J1644+2619: Its radio source has an inverted spectrum and was unresolved with VLBI at milli-arcsec resolution (Doi et al. 2007).

C. RADIO EMISSION AT LOW FREQUENCIES

In the above sections, evidence has been presented for the blazar-like properties of the (flat-spectrum) RL NLS1 galaxies, arguing for the presence of relativistic jets in them. As is known in normal radio quasars/galaxies, radio jets are associated with large scale radio lobes. These show diffuse radio emission with steep spectrum ($\alpha_r \simeq -0.77$; e.g. Kellermann 1966) which dominates the radio fluxes at low frequencies (several hundred MHz), via synchrotron radiation from optically thin plasma. Here we examine the radio properties of our sample at low frequencies, though the data are sparse in general, especially at around 100 MHz. The radio emission is searched for using the WENSS³¹ survey (Rengelink et al. 1997) at 327 MHz and using NED for surveys at lower frequencies. We find 15 objects with detections at 327 MHz, for which the 1.4 GHz to 327 MHz spectral indices $\alpha_{327M}^{1.4G}$ can be computed. We found a median $\alpha_{327M}^{1.4G}$ of -0.47 , and $\sim 47\%$ being steeper than -0.5 with the steepest as -0.83 , systematically steeper than the 1.4–5 GHz indices (3/11 are steep-spectrum). This means that the spectral indices steepen progressively towards low frequencies.

This behavior can well be explained by onset of a steep-spectrum component dominating the low frequencies, as commonly seen in normal RL AGN. Therefore, we postulate that in these objects there exists diffuse, optically thin emission, presumably from putative radio lobes that are powered by (relativistic) jets. If this is the case, those blazar-like (flat-spectrum) NLS1 AGN must possess the general core–halo morphology as seen in some blazars (e.g. Stannard & McIlwrath 1982; Browne & Perley 1986; Murphy et al. 1993). Given the compact morphology unresolved at the FIRST resolution (smaller than several tens kpc), there are two possibilities: either they have large scale extended emission—similar to the classical radio sources—but are viewed close to the radio axis, or are intrinsically compact, similar to CSS (compact steep-spectrum sources).

At the even lower frequency of 151 MHz, 4 objects have detected radio emission, all from the 6C/7C survey (e.g. Baldwin et al. 1985; Hales et al. 1988), namely J1047+4725 (1.81 Jy), J1305+5116 (0.32 Jy), J1443+4725 (0.5 Jy), and J1548+3511 (0.26 Jy). The 151–327 MHz indices are $\alpha_{151M}^{327M} = -0.28, -0.54, -0.62, +0.09$, respectively, and the average is -0.34 . These values are much flatter than the ‘canonical’ slope for diffuse optically thin emission from powerful sources (< -0.7 , e.g. Kellermann 1966), indicating that the spectra have already turned over, or started to turn over, in this frequency range due possibly to synchrotron self-absorption. In fact, the radio spectra of J1047+4725 and J1548+3511 show a clear trend of turnover at low frequencies. Considering the relatively large redshifts of the sample, the break frequencies in some sources could be as high as several hundred MHz in the source rest frame. Spectral turnovers at such frequencies are often seen in radio sources with small scale structure, such as in CSS (e.g. O’Dea & Baum 1997). The previously reported extreme NLS1/blazar hybrid object 2MASX J0324+3410 (Zhou et al. 2007) was also detected at 151 MHz (1.02 Jy); its radio slope also steepens significantly towards low frequencies, $\alpha_{151M}^{327M} = -0.44$ (compared to $\alpha_r \simeq 0.1$ at high frequencies, Neumann et al. 1994), suggesting the dominance of a steep-spectrum component at low frequencies.

D. EFFECTS OF REFRACTIVE INTERSTELLAR SCINTILLATION ON RADIO FLUX VARIATIONS

For compact radio sources (milli-arcsec scale), it is known that refractive interstellar scintillation (RISS), caused by inhomogeneities in the interstellar medium, may give rise to variations of radio flux densities (e.g. Rickett 1986, 1990; Blandford et al. 1986). RISS is wavelength-dependent and concerns predominantly the low frequency domain (below 1 GHz). At centimeter wavelengths, RISS mostly causes “flickering” at a level of a few percent, as was found in observations (Heeschen 1984) and is also consistent with the theoretical models (Rickett 1986; Blandford et al. 1986). Although variations at a larger amplitude were sometimes ascribed to RISS, they were limited to a small number of individual objects which ubiquitously are at low Galactic latitude ($< 10^\circ$) or even have a line-of-sight passing through the Galactic plane (Hjellming & Narayan 1986; Dennison et al. 1987). Monitoring observations of radio sources at frequencies above 1 GHz showed that both intrinsic and extrinsic (RISS) variability contribute to observed radio flux variations, with the former having timescales from months to years and the latter being faster and having rms amplitude typically $< 10\%$ (e.g. Lazio et al. 2001; Mitchell et al. 1994; Rickett et al. 2006).

In consideration of the large variation amplitudes in some of the objects of our sample and their high Galactic latitudes ($b=31^\circ - 51^\circ$, Table 3), we argue that the variations are predominantly intrinsic for the majority of the variables reported here, instead of arising from RISS. This is especially true for variations at 5 GHz or even higher frequencies. Furthermore, the brightness temperatures deduced from variability at 1.4 GHz are in the same range as those from 5 GHz (Table 3); this favours the intrinsic variability explanation, since RISS at 1.4 GHz is expected to be more pronounced than at 5 GHz due to its strong wavelength dependence. We therefore suggest that, although a non-negligible role played by RISS may not be ruled out in individual objects, especially those with relatively low amplitudes, our results for the sample as a whole should not be affected significantly.

E. ON THE X-RAY RADIATION MECHANISMS OF RADIO-LOUD NLS1 AGN

As one of the defining characteristics, the X-ray spectra of FSRQs are flat and extend from the soft to the hard X-ray band ($\Gamma \sim 1.6$ with a dispersion $\sigma = 0.1 - 0.2$; see e.g. Reeves & Turner 2000), which is interpreted as inverse Compton radiation from the relativistic jets and beamed. In most RLQs the contribution from a hot disk corona to the observed X-rays is negligible (in the hard X-ray band > 2 keV), except the steep-spectrum soft X-ray excess below 1 keV (e.g. Brinkmann et al. 1997; Yuan et al. 2000). From the data analysis above, we find a flat X-ray spectrum in J1633+4718 with $\Gamma = 1.37 \pm 0.49$, and possibly in

³¹ The Westerbork Northern Sky Survey

J0849+5108 with $\Gamma = 1.77^{+0.44}_{-0.60}$ (90% errors). Their concave optical to X-ray spectral shapes (see Figure 10) also agree with the inverse Compton scenario. For J0849+5108, the LBL-like SED and flat X-ray spectrum are well expected, based on its OVV/HPQ properties (see Appendix B). The prominent soft X-ray excess in J1633+4718 is also typical of normal NLS1 galaxies. It is puzzling, however, that J1633+4718 shows a broad band SED similar to that of HFSRQs, for which a steep X-ray spectrum and a convex α_{OX} -X-ray shape are generally expected. Future multi-waveband observations, especially in the X-ray band, are needed to confirm this peculiarity.

As discussed above, in the presence of an excess soft X-ray component, the effective spectral slopes derived from the hardness ratios are generally steep. The same is expected to be true for radio-loud NLS1 galaxies, as found in J1633+4718 above. For reference, FSRQs at redshifts ($z < 1$) are found to have $\Gamma \sim 2.1$ with a dispersion $\sigma \simeq 0.2$ in the ROSAT PSPC band (Brinkmann et al. 1997; Yuan 1998). Therefore, we expect that a few other objects, e.g. those having FSRQ/LBL-type SEDs yet steep Γ values in the PSPC bandpass (Figure 9) would similarly have a flat underlying X-ray continuum.

HBLs are commonly observed to show steep X-ray spectra (e.g. Worrall & Wilkes 1990; Brinkmann et al. 1996; Padovani & Giommi 1995; Wolter et al. 1998), which are intrinsically curved, progressively steepening with increasing photon energy (e.g. Perlman et al. 2005). Their X-ray emission is thought to be attributed to the high-energy tail of the synchrotron radiation, i.e. the synchrotron bump reaches the soft X-ray band (Ghisellini & Maraschi 1989). There are only a few flat-spectrum radio-loud NLS1 AGN studied previously with multi-waveband data and, interestingly, most of these show HBL/HFSRQ-like SEDs. For example, RXJ16290+4007, formerly classified as a FSRQ, was suggested by Padovani et al. (2002) to be the first confirmed HFSRQ for its modeled synchrotron peak at 2×10^{16} Hz and a steep soft X-ray spectrum $\Gamma \simeq 3$. 2MASX J0324+3410 also shows a continuum SED (square in Figure 9; Zhou et al. 2007) and a luminosity closely resembling that of Mkn 421, a well-known HBL; the object was claimed to be even marginally detected in TeV γ -rays (Falcone et al. 2004), as seen in some nearby bright HBL. In the X-ray band, 2MASX J0324+3410 also shows a steep spectrum ($\Gamma \simeq 2.2$, Zhou et al. 2007).

Most of the objects falling into the HBL locus in Figure 9 show steep X-ray spectra wherever available ($\Gamma = 2.0 - 3.3$). Meanwhile, their $\alpha_{\text{OX}} - \alpha_{\text{X}}$ SED shapes are either convex or close to a straight line (i.e. power-law; see Figure 10). Thus, based on the similarity with HBL in both the SED and X-ray spectral shape, we suggest their X-ray emission to be the high-energy tail of the synchrotron component, as in HBL. There are a few objects lying close to the border line, known as intermediate BL Lac objects (IBL) where the synchrotron peaks are around $10^{14} - 10^{15}$ Hz. In them, the high-energy synchrotron tail may still reach the soft X-rays, as found in a few IBL objects (e.g. Tagliaferri et al. 2000; Ferrero et al. 2006).

REFERENCES

- Abramowicz, M.A., Czerny, B., Lasota, J.P., Szuszkiewicz, E., 1988, *ApJ*, 332, 646
Adelman-McCarthy, J.K., et al. 2007, *ApJS*, 172, 634
Ai, Y. et al. 2008, in preparation
Arnaud, K.A., 1996, *Astronomical Data Analysis Software and Systems V*, eds. G. Jacoby and J. Barnes, p17, ASP Conf. Series volume 101
Arp, H., Sargent, W.L.W., Willis, A.G., Oosterbaan, C.E. 1979, *ApJ*, 230, 68
Bade, N., Fink, H.H., Engels, D., Voges, W., Hagen, H.-J., et al., 1995, *A&AS*, 110, 469
Bahcall, J.N., Kirhakos, S., Saxe, D.H., Schneider, D.P., 1997, *ApJ*, 479, 642
Baldwin, J.E., Boysen, R.C., Hales, S.E.G., Jennings, J.E., Waggett, P.C., et al. 1985, *MNRAS*, 217, 717
Ballantyne, D.R. 2007, *astro-ph/0707.3142*
Balmaverde, B. & Capetti, A. 2006, *A&A*, 447, 97
Baum, S.A., Zirbel, E.L., O'Dea, C. 1995, *ApJ*, 451, 88
Becker, R.H., White, R.L., & Edwards, A.L. 1991, *ApJS*, 75, 1
Becker, R.H., White, R.L. & Helfand, D.J. 1995, *ApJ*, 450, 559
Belloni, T., Hasinger, G., Izzo, C. 1994, *A&A*, 283, 1037
Best, P.N., Kauffmann, G., Heckman, T.M., Brinchmann, J., Charlot, S., et al. 2005, *MNRAS*, 362, 25
Bian, W. & Zhao, Y. 2004, *MNRAS*, 352, 823
Blackburn, J.K. 1995, in *ASP Conf. Ser.*, Vol. 77, *Astronomical Data Analysis Software and Systems IV*, ed. R.A. Shaw, H.E. Payne, and J.J.E. Hayes (San Francisco: ASP), 367 (<http://heasarc.gsfc.nasa.gov/f-tools>)
Blandford, R.D. & Rees, M.J. 1978, in *Pittsburgh Conference on BL Lac Objects*, ed. A.M. Wolfe (Pittsburgh: Univ. Pittsburgh Dept. Phys. Astron.), p328
Blandford, R.D. & Königl, A. 1979, *ApJ*, 232, 34
Blandford, R.D., Narayan, R., Romani, R.W. 1986, *ApJ*, 301, L53
Blandford, R.D. 2000, *Roy. Soc. of London Phil. Tr. A.*, vol. 358, issue 1767, p.811
Blundell, K.M., Beasley, A.J., & Bicknell, G.V. 2003, *ApJ*, 591, L103
Blundell, K.M. & Rawlings, S. 2001, *ApJ*, 562, L5
Boller, Th., Meurs, E.I.A., Brinkmann, W., Fink, H., Zimmermann, U., Adorf, H.-M. 1992, *A&A*, 261, 57
Boller, T., Brandt, W.N., & Fink, H. 1996, *A&A*, 305, 53
Boroson, T.A. & Green, R.F. 1992, *ApJS*, 80, 109
Boroson, T.A. 2002, *ApJ*, 565, 78
Brinkmann, W., & Siebert, J. 1994, *A&A*, 285, 812
Brinkmann, W., Siebert, J., Kollgaard, R.I., Thomas, H.-C. 1996, *A&A*, 313, 356
Brinkmann, W., Yuan, W., Siebert, J. 1997, *A&A*, 319, 413
Browne, I.W.A., & Perley, R.A. 1986, *MNRAS*, 222, 149
Brunthaler, A., et al. 2005, *A&A*, 435, 497
Bruzual, G. & Charlot, S. 2003, *MNRAS*, 344, 1000
Cao, X. 2003, *ApJ*, 599, 147
Capetti, A. & Balmaverde, B. 2006, *A&A*, 453, 27
Cash, W. 1979, *ApJ*, 228, 939
Celotti, A., Padovani, P., Ghisellini, G. 1997, *MNRAS*, 286, 415
Celotti, A. & Blandford, R.D. 2001, in 'Black Holes in Binaries and Galactic Nuclei: Diagnostics, Demography and Formation', (Eds) L. Kaper, E.P.J. van den Heuvel, & P.A. Woudt. Springer-Verlag, 2001, p.206
Celotti, A. 2002, in 'Relativistic Flows in Astrophysics', (Eds) A.W. Guthmann et al., *LNP* 589, p.88
Collin, S., & Kawaguchi, T. 2004, *A&A*, 426, 797
Collin, S., Kawaguchi, T., Peterson, B.M., Vestergaard, M. 2006, *A&A*, 456, 75
Condon, J.J., Cotton, W.D., Greisen, E.W., Yin, Q.F., Perley, R.A., et al. 1998, *AJ*, 115, 1693
Condon, J.J., Yin, Q.F., Thuan, T.X., Boller, Th. 1998, *AJ*, 116, 2682
Constantin, A. & Shields, J.C. 2003, *PASP*, 115, 592
Corbett, E.A., Robinson, A., Axon, D.J., and Hough, J.H. 2000, *MNRAS*, 311, 485
Dallacasa, D., Bondi, M., Alef, W., Mantovani, F. 1998, *A&AS*, 129, 219
Decarli R., Dotti M., Fontana M., Haardt F., 2008, *MNRAS*, in press (*astro-ph/0801.4560*)
Dennison, B., et al. 1987, *ApJ*, 313, 141
Dietrich, M., Hamann, F., Appenzeller, I. & Vestergaard, M. 2003, *ApJ*, 596, 817
Doi, A., Nagai, H., Asada, K., Kamenoi, S., Wajima, K., & Inoue, M. 2006, *PASJ*, 58, 777
Doi, A., Fukusawa, K., Inoue, M., Wajima, K., Nagai, H., et al. 2007, *PASJ*, accepted (*astro-ph/0704.3836*)
Dong, X.-B., Zhou, H.-Y., Wang, T.-G., Wang, J.-X., Li, C., & Zhou, Y.-Y. 2005, *ApJ*, 620, 629
Dong, X.-B., Wang, T.-G., Wang, J., Yuan, W., Zhou, H.-Y., Dai, H.-F., & Zhang, K. 2008, *MNRAS*, 383, 581
Dunlop, J.S., McLure, R.J., Kukula, M.J., Baum, S.A., O'Dea, C.P., Hughes, D.H., 2003, *MNRAS*, 340, 1095
Falcke, H., Sherwood, W., & Patnaik, A.R. 1996a, *ApJ*, 471, 106
Falcke, H., Patnaik, A.R., & Sherwood, W. 1996b, *ApJ*, 473, L13
Falcone, A.D., et al. 2004, *ApJ*, 613, 710
Fanaroff, B.L. & Riley, J.M. 1974, *MNRAS*, 167, 31

- Fanti, C., Fanti, R., Ficarra, A., Gregorini, L., Mantovani, F., Padrielli, L. 1983, *A&A*, 118, 171
- Fender, R.P., Bellone, T.M., Gallo, E. 2004, *MNRAS*, 2004, 355, 1105
- Ferrero, E., Wagner, S.J., Emmanoulopoulos, D., Ostorero, L. 2006, *A&A*, 457, 133
- Floyd, D.J.E., et al. 2004, *MNRAS*, 355, 196
- Fossati, G., Maraschi, L., Celotti, A., Comastri, A., Ghisellini, G. 1998, *MNRAS*, 299, 433
- Francis, P.J. 1996, *Publications of the Astronomical Society of Australia*, 13, 212
- Francis, P.J. 1991, *ApJ*, 373, 465
- Fugmann, W. 1988, *A&A*, 205, 86
- Gallo, L.C., et al. 2006, *MNRAS*, 370, 245
- Ghisellini, G. & Maraschi, L., 1989, *ApJ*, 340, 181
- Ghisellini, G., Padovani, P., Celotti, A., Maraschi, L. 1993, *ApJ*, 407, 65
- Ghisellini, G., Celotti, A., Fossati, G., Maraschi, L., Comastri, A. 1998, *MNRAS*, 301, 451
- Ghosh, K.K. & Punsly, B., 2007, astro-ph/0704.2407
- Goodrich, R.W. 1989, *ApJ*, 342, 22
- Gopal-Krishna, Krishnamohan, S., Singal, A.K. 1984, *A&A*, 140, L19
- Gopal-Krishna & Wiita, P.J., 2002, *NewAR*, 46, 357
- Greene, J.E. & Ho, L.C. 2005, *ApJ*, 630, 122
- Greene, J.E., Ho, L.C., Ulvestad, J.S. 2006, *ApJ*, 636, 56
- Greene, J.E. & Ho, L.C. 2007, *ApJ*, 670, 92
- Griffith, M.R., Wright, A.E., Burke, B.F., Ekers, R.D. 1995, *ApJS*, 97, 347
- Grupe, D., Beuermann, K., Thomas, H.-C., Mannheim, K., & Fink, H.H. 1998, *A&A*, 330, 25
- Grupe, D., Beuermann, K., Mannheim, K., & Thomas, H.-C. 1999, *A&A*, 350, 805
- Grupe, D., Wills, B.J., Leighly, K.M., & Meusinger, H. 2004, *AJ*, 127, 156
- Hales, S.E.G., Baldwin, J.E., Warner, P.J. 1988, *MNRAS*, 234, 919
- Heeschen, D.S. 1984, *AJ*, 89, 1111
- Hjellming, R.M. & Narayan, R. 1986, *ApJ*, 310, 768
- Ho, L.C., & Peng, C.Y. 2001, *ApJ*, 555, 650
- Ho, L.C., & Ulvestad, J.S. 2001, *ApJS*, 133, 77
- Ho, L.C. 2002, *ApJ*, 564, 120
- Isobe, T., Feigelson, E.D., & Nelson, P.I., 1986, *ApJ*, 306, 490
- Isobe, T. & Feigelson, E.D. 1990, "ASURV", *Bull. Amer. Astro. Society*, 22, 917
- Ivezić, Ž., et al. 2002, *AJ*, 124, 2364
- Jackson, N., Browne, I.W.A. 1991, *MNRAS*, 250, 422
- Jarvis, M.J., & McLure, R.J. 2002, *MNRAS*, 336, L38
- Jarvis, M.J. & McLure, R.J. 2006, *MNRAS*, 369, 182
- Jiang, D.R., Cao, X., Hong, X. 1998, *ApJ*, 494, 139
- Jones, T.W. & Burbidge, G.R. 1973, *ApJ*, 186, 791
- Kaspi, S., Maoz, D., Netzer, H., Peterson, B.M., Vestergaard, M., Jannuzi, B.T. 2005, *ApJ*, 629, 61
- Kaspi, S., Smith, P.S., Netzer, H., Maoz, D., Jannuzi, B.T., & Givon, U. 2000, *ApJ*, 533, 631
- Kellermann, K.I. 1966, *ApJ*, 146, 621
- Kellermann, K.I., & Pauliny-Toth, I.I.K. 1969, *ApJ*, 155, L71
- Kellermann, K.I., Sramek, R., Schmidt, M., Shaffer, D.B., Green, R. 1989, *AJ*, 98, 1195
- Kellermann, K.I., Sramek, R.A., Schmidt, M., Green, R.F., Shaffer, D.B. 1994, *AJ*, 108, 1163
- Komossa, S., & Meersweinchen, J. 2000, *A&A*, 354, 411
- Komossa, S., Voges, W., Adorf, H.M., Xu, D., Mathur, S., Anderson, S. 2006, *ApJ*, 639, 710
- Komossa, S., Voges, W., Xu, D., Mathur, S., Adorf, H.M., Lemson, G., Duschl, W., Grupe, D. 2006, *AJ*, 132, 531
- Komossa, S., Xu, D., Zhou, H., Binette, L., Storchi-Bergmann, T. 2007, *ApJ*, submitted
- Komossa, S. 2008, in: *The nuclear region, host galaxy and environment of AGN*, E. Benitez, I. Cruz-Gonzales, Y. Krongold (eds), *RevMexAA*, in press (arXiv:0710.3326v1)
- Körting, E.G., Jester, S., Fender, R.P. 2006, *MNRAS*, 372, 1366
- Krolik, J.H. 1999, *Active Galactic Nuclei* (Princeton Univ. Press)
- Lacy, M., Laurent-Muehleisen, S.A., Ridgway, S.E., Becker, R.H., White, R.L., 2001, *ApJ*, 551, L17
- Laing, R.A., Riley, J.M., Longair, M.S., 1983, *MNRAS*, 204, 151
- Langston, G.I., Heflin, M.B., Conner, S.R., Lehar, J., Carrilli, Ch.L., Burke, B.F. 1990, *ApJS*, 72, 621
- Laor, A. 2000, *ApJ*, 543, L111
- Laor, A. 2003, arXiv:astro-ph/0312417v2
- Lazio, T.J.W., Waltman, E.B., Ghigo, F.D., Fiedler, R.L., Foster, R.S., Johnston, K.J. 2001, *ApJS*, 136, 265
- Leighly, K.M. 1999, *ApJS*, 125, 297
- Leighly, K.M. 2000, *New Astronomy Review*, 44, 395
- Liu, Y., Jiang, D.R., Gu, M.F. 2006, *ApJ*, 637, 669
- Lu, H., Zhou, H., Wang, J., Wang, T., Dong, X., Zhuang, Z., & Li, C. 2006, *AJ*, 131, 790
- Lu, Y., Wang, T., Zhou, H., Wu, J., 2007, *AJ*, 133, 1615
- Maccacaro, T., Gioia, I.M., Wolter, A., Zamorani, G., Stocke, J.T. 1988, *ApJ*, 326, 680
- Maccarone, T.J., Gallo, E., Fender, R. 2003, *MNRAS*, 345, L19
- Magorrian, J., Tremaine, S., Richstone, D., Bender, R., Bower, G. et al., 1998, *AJ*, 115, 2285
- Maraschi, L. & Tavecchio, F. 2003, *ApJ*, 593, 667
- Martin, D.C., et al. 2005, *ApJ*, 619, L1
- McLure, R.J., & Jarvis, M.J. 2004, *MNRAS*, 353, L45
- Meier, D.L. 2003, *New Astronomy Reviews*, 47, 667
- Metcalf, R.B. & Magliocchetti, M. 2006, *MNRAS*, 365, 101
- Miller, L., Peacock, J.A., Mead, A.R.G., 1990, *MNRAS*, 244, 207
- Mineshige, S., Kawaguchi, T., Takeuchi, M., Hayashida, K., 2000, *PASJ*, 52, 499
- Mitchell, K.J., Dennison, B., Condon, J.J., Altschuler, D.R., Payne, H.E., et al. 1994, *ApJS*, 93, 441
- Moore, R.L., & Stockman, H.S., 1981, *ApJ*, 243, 60
- Moran, E.C., Halpern, J.P., Helfand, D.J. 1996, *ApJS*, 106, 341
- Moran, E.C. 2000, *NewA Rev.*, 44, 527
- Murphy, D.W., Browne, I.W.A., & Perley, R. A. 1993, *MNRAS*, 264, 298
- Nousek, J.A., & Shue, D.R. 1989, *ApJ*, 342, 1207
- Netzer, H. 1990, in *Active Galactic Nuclei*, ed. T.J.-L. Courvoisier & M. Mayor (Berlin: Springer), 137
- Neumann, M., Reich, W., Furst, E., Brinkmann, W., Reich, P., et al. 1994, *A&AS*, 106, 303
- O'Dea, C.P. & Baum, S.A. 1997, *AJ*, 113, 148
- O'Dea, C.P. 1998, *PASP*, 110, 493
- O'Dowd, M. & Urry, C.M. 2005, *ApJ*, 627, 97
- Ohsuga, K., Mori, M., Nakamoto, T., Mineshige, S. 2005, *ApJ*, 628, 368
- Onken, C.A., Ferrarese, L., Merritt, D., Peterson, B.M., Pogge, R.W., et al. 2004, *ApJ*, 615, 645
- Örndahl, E. & Rönnback, J. 2005, *A&A*, 443, 61
- Oshlack, A.Y.K.N., Webster, R.L., & Whiting, M.T. 2001, *ApJ*, 558, 578
- Oshlack, A.Y.K.N., Webster, R.L., & Whiting, M.T. 2001, *ApJ*, 576, 81
- Osterbrock, D.E., & Pogge, R.W. 1985, *ApJ*, 297, 166
- Padovani, P. & Giommi, P. 1995, *ApJ*, 444, 567
- Padovani, P. & Giommi, P., 1996, *MNRAS*, 279, 526
- Padovani, P., Giommi, P., Fiore, F. 1997a, *MNRAS*, 284, 569
- Padovani, P., Giommi, P., Fiore, F. 1997b, in *Mem. Soc. Astron. Italiana*, 68, 147
- Padovani, P., Costamante, L., Ghisellini, G., Giommi, P., Perlman, E. 2002, *ApJ*, 581, 895
- Padovani, P., Perlman, E., Landt, H., Giommi, P., Perri, M. 2003, *ApJ*, 588, 128
- Pei, Y.C. 1992, *ApJ*, 395, 130
- Perlman, E.S., Padovani, P., Giommi, P., Sambruna, R., Jones, L.R., et al. 1998, *AJ*, 115, 1253
- Perlman, E.S., Madejski, G., Georgopoulos, M., Andersson, K., Daugherty, T., et al. 2005, *ApJ*, 625, 727
- Puchnarewicz, E.M., et al. 1992, *MNRAS*, 256, 589
- Punsly, B. 2007, *ApJ*, 657, L9
- Rawlings, S. & Saunders, R. 1991, *Nature*, 349, 138
- Readhead, A.C.S. 1994, *ApJ*, 426, 51
- Reeves, J.N. & Turner, M.J.L. 2000, *MNRAS*, 316, 234
- Reich, W., Fürst, E., Reich, P., Kothes, R., Brinkmann, W., & Siebert, J., 2000, *A&A*, 363, 141
- Rengelink, R.B., Tang, Y., de Bruyn, A.G., Miley, G.K., Bremer, M.N., et al. 1997, *A&AS*, 124, 259
- Rickett, B.J. 1986, *ApJ*, 307, 564
- Rickett, B.J. 1990, *ARA&A*, 28, 561
- Rickett, B.J., Lazio, T.J.W., Ghigo, F.D. 2006, *ApJS*, 165, 439
- Sambruna, R.M., Maraschi, L., Urry, C.M. 1996, *ApJ*, 463, 444
- Schartel, N., et al. 1996, *MNRAS*, 283, 1015
- Schwöpe, A.D., et al. 2000, *Astron. Nachr.*, 321, 1
- Siebert, J. 1996, Ph.D. Thesis, Techn. Univ. München
- Sikora, M., Stawarz, L., Lasota, J.-P. 2007, *ApJ*, 2007, 658, 815
- Singal, A.K. 1993, *MNRAS*, 263, 139
- Sitko, M.L., Rudnick, L., Jones, T.W., Schmidt, G.D. 1984, *PASP*, 96, 402
- Smith, D.A., Herter, T., & Haynes, M.P. 1998, *ApJ*, 494, 150
- Smith, J.E., Robinson, A., Alexander, D.M., Young, S., Axon, D.J., Corbett, E.A. 2004, *MNRAS*, 350, 140
- Stalin, C.S., Gupta, A.C., Gopal-Krishna, Wiita, P.J., Sagar, R. 2005, *MNRAS*, 356, 607
- Stannard, D. & McIlwrath, B.K. 1982, *Nature*, 298, 140
- Stein, W.A., Odell, S.L., Strittmatter, P.A. 1976, *ARA&A*, 14, 173

- Stepanian, J.A. 2003, *ApJ*, 588, 746
- Stocke, J.T., Morris, S.L., Weymann, R.J., Foltz, C.B. 1992, *ApJ*, 396, 487
- Stickel, M., Fred, J.W., Kühr, H. 1989, *A&A*, 224, L27
- Sulentic, J.W., Zwitter, T., Marziani, P., & Dultzin-Hacyan, D. 2000, *ApJ*, 536, L5
- Sulentic, J.W., Zamfir, S., Marziani, P., Bachev, R., Calvani, M., Dultzin-Hacyan, D. 2003, *ApJ*, 597, L17
- Tadhunter, C., Wills, K., Morganti, R., Oosterloo, T., Dickson, R. 2001, *MNRAS*, 327, 227
- Tagliaferri, G., Ghisellini, G., Giommi, P., Chiappetti, L., Maraschi, L., Celotti, A., et al. 2000, *A&A*, 354, 431
- Tinti, S., Dallacasa, D., De Zotti, G., Celotti, A., Stanghellini, C. 2005, *A&A*, 432, 31
- Ulvestad, J.S., Antonucci, R.R.J., & Goodrich, R.W. 1995, *AJ*, 109, 81
- Urry, C.M. & Padovani, P. 1995, *PASP*, 107, 803
- Vermeulen, R.C., et al. 1995, *ApJ*, 452, L5
- Véron-Cetty, M.-P., Véron, P., & Gonçalves, A.C. 2001, *A&A*, 372, 730
- Véron-Cetty, M.-P., Joly, M., & Véron, P. 2004, *A&A*, 417, 515
- Vestergaard, M. 2004, *ApJ*, 601, 676
- Vestergaard, M., Wilkes, B.J., Barthel, P.D. 2000, *ApJ*, 538, L103
- Vestergaard, M. & Peterson, B.M. 2006, *ApJ*, 641, 689
- Voges, W., et al. 1999, *A&A*, 349, 389
- Warner, C., Hamann, F., Dietrich, M. 2004, *ApJ*, 608, 136
- Wandel, A., Peterson, B.M., & Malkan, M.A. 1999, *ApJ*, 526, 579
- Wang, J.-M., & Netzer, H. 2003, *A&A*, 398, 927
- Wang, T., Brinkmann, W. & Bergeron, J. 1996, *A&A*, 309, 81
- Wang, T.G., Matsuoka, M., Kubo, H., Mihara, T., & Negoro, H. 2001, *ApJ*, 554, 233
- Wang, T.G., Dong, X.B., Zhou, H.Y., & Wang, J. X. 2005, *ApJ*, 622, L101
- Wang, T.-G., Zhou, H.-Y., Wang, J.-X., Lu, Y.-J., Lu, Y. 2006, *ApJ*, 645, 856
- Whalen, D.J., Laurent-Muehleisen, S.A., Moran, E.C., Becker, R.H. 2006, *AJ*, 131, 1948
- White, N.E., Giommi, P., & Angelini, L. 1995, WGA ROSAT source catalog
- Whiting, M.T., Webster, R.L., & Francis, P.J. 2001, *MNRAS*, 323, 718
- Williams, R.J., Pogge, R.W., Mathur, S. 2002, *AJ*, 124, 3042
- Willis, A.G., Oosterbaan, C.E., de Ruiter H.R. 1976, *A&AS*, 25, 543
- Wills, B.J. & Browne, I.W.A. 1986, *ApJ*, 302, 56
- Wisozki, L., & Bade, N. 1997, *A&A*, 320, 395
- Wolter, A., Comastri, A., Ghisellini, G., Giommi, P., Guainazzi, M., et al. 1998, *A&A*, 335, 899
- Woo, J.-H. & Urry, C.M. 2002, *ApJ*, 581, L5
- Worrall, D.M. & Wilkes, B.J. 1990, *ApJ*, 360, 396
- Wyder, T.K., et al. 2005, *ApJ*, 619, L15
- Yuan, W. 1998, Ph.D. Thesis, Techn. Univ. München
- Yuan, W., Brinkmann, W., Siebert, J., Voges, W. 1998, *A&A*, 330, 108
- Yuan, W., Brinkmann, W., Gliozzi, M., Zhao, Y., Matsuoka, M. 2000, *A&A*, 362, 19
- Zhou, H.-Y., & Wang, T.-G. 2002, *Chinese Journal of Astron. & Astrophy.*, 2, 501
- Zhou, H.-Y., Wang, T.-G., Dong, X.-B., Zhou, Y.-Y., & Li, C. 2003, *ApJ*, 584, 147
- Zhou, H.-Y., Wang, T.-G., Dong, X.-B., Li, C., & Zhang, X.-G. 2005, *Chinese Journal of Astron. & Astrophy.*, 5, 41
- Zhou, H., Wang, T., Yuan, W., Lu, H., Dong, X., Wang, J., Lu, Y. 2006, *ApJS*, 166, 128
- Zhou, H., Wang, T., Yuan, W., Shan, H., Komossa, S., et al. 2007, *ApJ*, 658, L13

TABLE 1
SAMPLE OF VERY RADIO-LOUD NLS1 GALAXIES

No.	name	SDSS name	Z	u mag	g mag	M_B mag	$f_{20\text{cm}}$ mJy	$f_{6\text{cm}}$ mJy	$f_{92\text{cm}}$ mJy	α_r	$\log R$	λf_{5100}^a	α_λ	$\text{fwhm}(\text{H}\beta)$ km s^{-1}	$f(\text{H}\beta)^b$	$f(\text{OIII})^b$	R_{4570}	m_{NUV} mag	m_{FUV} mag
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)
1	J0814+5609	J081432.11+560956.6	0.509	18.05	17.89	-24.1	69/60	43	72	-0.33	2.53	458	-2.43	2164± 59	870±16	54± 7	1.04		
2	J0849+5108	J084957.98+510829.0	0.583	19.80	18.85	-23.1	344/266	161	202	~ 0 ^c	3.16	187	-0.20	1811±191	135±12	42± 3	0.23		
3	J0850+4626	J085001.17+462600.5	0.523	19.67	19.02	-22.8	21/16	...	51	...	2.23	409	-0.86	1251± 61	255±10	133± 5	1.48	20.80	21.29
4	J0902+0443	J090227.16+044309.6	0.532	19.38	18.92	-23.0	153/157	106	...	-0.30	3.02	520	-0.53	2089±205	341±20	192± 6	1.10	20.28	
5	J0948+0022	J094857.32+002225.5	0.584	18.59	18.36	-23.9	108/70	295	...	0.82	2.55	510	-1.87	1432± 87	341±12	34± 4	1.22	18.61	19.04
6	J0953+2836	J095317.09+283601.5	0.657	19.19	18.94	-23.6	45/43	...	...	...	2.71	240	-1.73	2162±201	215±14	52± 5	1.23		
7	J1031+4234	J103123.73+423439.3	0.376	19.64	19.36	-21.6	17/19	...	44	...	2.34	284	-0.55	1642± 75	379±14	131± 8	1.07		
8	J1037+0036	J103727.45+003635.6	0.595	20.04	19.32	-22.8	27/28	...	...	...	2.66	225	-1.71	1357±107	173± 9	25± 4	1.58	21.25	22.99
9	J1047+4725	J104732.68+472532.1	0.798	19.16	18.95	-24.0	734/789	404	1463	-0.51	3.87	293	-1.30	2153±267	421±32	325± 8	0.66		
10	J1110+3653	J111005.03+365336.3	0.630	21.13	20.59	-21.8	19/23	...	45	...	2.97	74	-1.28	1300±339	93±20	32± 4	0.88	21.35	21.85
11	J1138+3653	J113824.54+365327.1	0.356	20.49	19.52	-21.2	13/12	...	26	...	2.34	378	...	1364± 99	393±20	72±10	0.69		
12	J1146+3236	J114654.28+323652.3	0.465	19.08	18.62	-23.0	15/17	...	26	...	2.11	427	-1.54	2081±183	632±39	149± 6	0.78		
13	J1238+3942	J123852.12+394227.8	0.622	19.89	19.55	-22.8	10/10	...	...	...	2.23	175	-1.19	910±200	103±18	68± 4	1.05		
14	J1246+0238	J124634.65+023809.0	0.362	18.52	18.31	-22.7	37/36	...	...	...	2.38	601	-1.89	1425± 68	527±12	115± 5	0.84	19.01	19.58
15	J1305+5116	J130522.75+511640.3	0.785	17.45	17.28	-25.6	84/87	46	211	-0.50	2.34	1264	-1.68	1925± 53	1982±43	487±13	0.65	17.45	18.14
16	J1435+3131	J143509.49+313147.8	0.501	19.87	19.43	-22.3	39/43	...	137	...	2.87	306	-1.70	1719±253	346±36	137± 6	1.32		
17	J1443+4725	J144318.56+472556.7	0.703	18.20	18.11	-24.6	165/166	72	309	-0.67	3.07	534	-2.06	1848±113	337±18	105±13	1.45	18.09	18.57
18	J1505+0326	J150506.48+032630.8	0.408	18.78	18.52	-22.7	365/395	859	...	0.66	3.19	389	-0.83	1082±113	95± 8	106± 4	1.53		
19	J1548+3511	J154817.92+351128.0	0.478	18.18	17.91	-23.9	141/141	107	279	-0.22	2.84	545	-2.32	2035± 52	721±12	309± 7	1.24	18.41	18.69
20	J1633+4718	J163323.58+471859.0	0.116	17.94	17.38	-20.9	63/69	35	107	-0.30	2.22	1868	-1.70	909± 43	902±21	919± 8	1.02	18.20	19.48
21	J1634+4809	J163401.94+480940.2	0.494	20.18	19.53	-22.2	8/14	...	16	...	2.31	257	-1.60	1609± 79	287±11	87± 3	0.98	21.03	22.30
22	J1644+2619	J164442.53+261913.2	0.144	18.03	18.02	-20.8	88/128	99	...	-0.07	2.65	1788	-1.00	1507± 42	1108±22	120± 4	0.75	18.50	18.65
23	J1722+5654	J172206.03+565451.6	0.425	18.66	18.31	-23.1	37/43	...	119	...	2.37	544	-1.41	1385± 36	542±11	323± 4	0.58		

NOTE. — Col.(2): abbreviated name of object used in this paper; Col.(3): SDSS name; Col.(4): redshift; Col.(5): SDSS u band PSF magnitude corrected for Galactic extinction; Col.(6): as col.(5), but for the g band; Col.(7): derived B band absolute magnitude; Col.(8): FIRST and NVSS radio flux densities at 20 cm (FIRST/NVSS); Col.(9): flux density at 6 cm; Col.(10): flux density at 92 cm given by WNESS; Col.(11): radio spectral slope between 6 cm and 20 cm; Col.(12): logarithm of radio-loudness defined as $\equiv f_\nu(1.4\text{GHz})/f_\nu(4400)$; Col.(13): flux λf at 5100 Å in the rest frame; Col.(14): optical power-law slope ($S(\lambda) \propto \lambda^{\alpha_\lambda}$); Col.(15): line width of H β in FWHM; Col.(16): line flux of the broad component of H β ; Col.(17): line flux of [O III]; Col.(18): FeII-to-H β flux ratio (see text); Col.(19): near-UV magnitude given by GALEX corrected for Galactic extinction; Col.(20): the same as col.(19) but for the far-UV band.

^aRest frame flux λf in units of $10^{-15} \text{ erg s}^{-1} \text{ cm}^{-2}$

^bRest frame line flux in units of $10^{-17} \text{ erg s}^{-1} \text{ cm}^{-2}$

^cGiven the strong variation of the 1.4 GHz flux, we adopt a representing radio slope $\alpha_r \sim 0$ as suggested by the radio data from Arp et al. (1979).

TABLE 2
X-RAY DATA AND ANALYSIS RESULTS

name	X-ray source	obs.	$T_{\text{expo}}(\text{s})$ sec	$N_{\text{H,Gal}}$ 10^{20} cm^{-2}	C.Rate 10^{-2} c/s	flux ^a	Γ	kT_{bb} eV	remark
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
J0814+5609	J081432.9+561008	RS	345	4.49	5.40	1.71	$2.57^{+0.40}_{-0.46}$		BSC
—	J081432.11+560956.6	RP	7343	4.49	2.04	0.39	3.36 ± 0.12		WGA
J0849+5108	J084957.98+510829.0	RP	4472	3.00	1.19	0.27	$1.77^{+0.32}_{-0.39}$		fit
J0850+4626	no detection		469	2.64	<4.51	<0.98	...		
J0902+0443	no detection		362	3.11	<5.72	<1.45	...		
J0948+0022	J094856.9+002235	RS	447	5.22	4.10	1.16	2.26 ± 0.64		FSC
J0953+2836	no detection		449	1.28	<5.52	<0.66	...		
J1031+4234	no detection		498	0.99	<6.00	<0.60	...		
J1037+0036	J103727.45+003635.6	RP	10475	4.87	0.84	0.16	$2.43^{+0.43}_{-0.49}$		
—	J103727.45+003635.6	RP	6158	4.87	2.56	0.51	3.23 ± 0.29		
J1047+4725	J104731.4+472528	RS	509	1.30	2.13	0.26	...		FSC
J1110+3653	no detection		315	1.69	<7.22	<1.06	...		
J1138+3653	no detection		327	1.75	<6.16	<0.94	...		
J1146+3236	J114654.0+323653	RS	452	1.42	1.67	0.21	$2.37^{+0.81}_{-0.60}$		FSC
J1238+3942	J123852.1+394253	RS	508	1.46	5.08	0.65	$2.45^{+0.32}_{-0.28}$		BSC
J1246+0238	no detection		160	2.01	<11.6	<1.98	...		
J1305+5116	no detection		499	0.98	<5.82	<0.58	...		
J1435+3131	J143509.2+313203	RS	382	1.14	10.50	1.15	...		BSC
J1443+4725	J144319.8+472541	RS	587	1.47	1.92	0.25	...		FSC
J1505+0326	no detection		417	3.89	<6.90	<2.14	...		
J1548+3511	J154817.6+351130	RS	241	2.28	5.49	1.05	...		FSC
J1633+4718	J163323.58+471859.0	RP	3748	1.79	26.0	5.44	1.37 ± 0.30	$32.5^{+3.9}_{-3.7}$	fit
J1633+4718	J163323.3+471848	RS	909	1.79	20.0	4.72	$1.47^{+0.77}_{-0.92}$	30^{+12}_{-10}	fit
J1634+4809	J163400.1+480915	RS	943	1.64	1.40	0.20	...		FSC
J1644+2619	J164443.2+261909	RS	518	5.12	10.6	2.60	$2.03^{+0.25}_{-0.28}$		BSC
—	as target	Ch	2946	5.12	18.0 ^b	1.05 ^b	2.19 ± 0.17		fit
J1722+5654	J172205.0+565450	RS	1415	2.10	8.11	1.33	2.40 ± 0.15		BSC

NOTE. — Col.(1): object name; Col.(2): name of X-ray source; Col.(3): RS – the RASS, RP – ROSAT pointed observation, Ch – Chandra observation; Col.(4): effective exposure time in units of seconds (vegneting corrected); Col.(5): Galactic absorption column density; Col.(6): X-ray count rate as detected with the ROSAT PSPC (except one of J1644+2619 with the Chandra ACIS-S; Col.(7): X-ray flux density corrected for Galactic absorption (0.1–2.4 keV for ROSAT and 0.3–5 keV band for Chandra data); Col.(8): power-law photon index; Col.(9): temperature of an additional soft X-ray blackbody component; Col.(10): remarks on how the X-ray data analysis is performed; fit – by spectral fitting; for the rest, the fluxes and Γ are derived using count rates and hardness ratio (see text for details) from RASS bright source catalogue (BSC), RASS faint source catalogue (FSC), and ROSAT source catalogues from pointed observations (RXP) and its WGA version (WGA), respectively.

^ain units of $10^{-12} \text{ erg s}^{-1} \text{ cm}^{-2}$

^bin the 0.3–5 keV band for the Chandra ACIS-S detector

TABLE 3
RADIO FLUX VARIATIONS AND INFERRED BRIGHTNESS TEMPERATURES

name	b deg.	band GHz	ΔS mJy	$\Delta S/\langle S \rangle$ %	Δt days	$\log T_B$ K	$\delta_{\min}$	reference
J0814+5609	33.7	1.4	9	14	1273	12.0	-	2
J0849+5108	39.1	1.4	78	25	1257	13.0	2.2	2,6
.....		1.4	230	75	2191	13.0	2.2	7,8
J0850+4626	39.3	1.4	5	27	1223	11.8	-	2,6
J0902+0443	31.2	4.85	10	9	1125	11.1	-	1,3
J0948+0022	38.7	1.4	38	43	1282	12.7	1.7	2,6
J1505+0326	50.3	4.85	527	72	1125	12.6	1.5	1,3
.....		8.3	146	43	64	14.0	4.7	9
J1548+3511	51.6	4.85	38	44	207	13.0	2.2	1,4
J1633+4718	42.6	4.85	24	55	1370	9.9	-	1,5

NOTE. — band – observing frequency; b – Galactic latitude; ΔS – absolute flux change between two epoches; $\Delta S/\langle S \rangle$ – fractional amplitudes where $\langle S \rangle$ is the average of the two fluxes; Δt – time span of the two epoches (for an observation whose epoch is available accurately only to month, the maximum length of dates is used for conservative estimation); T_B – brightness temperature (logarithm); $\delta_{\min}$ – minimum Doppler factor of beaming assuming an intrinsic maximum T_B as the inverse Compton limit 10^{12} K. Reference codes: (1) Becker et al. (1991) (2) Becker et al. (1995) (3) Griffith et al. (1995) (4) Langston et al. (1990) (5) Neumann et al. (1994) (6) Condon et al. (1998a) (7) Willis et al. (1976) (8) Arp et al. (1979) (9) Goddard Geodetic VLBI GROUP’s auxiliary web pages <http://lacerta.gsfc.nasa.gov/vlbi/images/>

TABLE 4
BROAD-BAND EFFECTIVE SPECTRAL INDICES^a AND
BLACK HOLE MASSES^b

name	α_{ro}	α_{rx}	α_{ox}	$\log(M_{\text{BH}}/M_{\odot})$
J0814+5609	0.44	0.87	1.75	8.0
J0849+5108	0.65	0.84	1.23	7.4
J0850+4626	0.44	>0.68	>1.17	7.2
J0902+0443	0.60	>0.78	>1.16	7.7
J0948+0022	0.57	0.78	1.22	7.5
J0953+2836	0.48	>0.74	>1.29	7.8
J1031+4234	0.44	>0.71	>1.27	7.3
J1037+0036	0.50	0.79	1.39	7.3
J1047+4725	0.70	0.94	1.45	8.1
J1110+3653	0.56	>0.68	>0.92	7.1
J1138+3653	0.47	>0.66	>1.07	7.1
J1146+3236	0.39	0.74	1.47	7.8
J1238+3942	0.41	0.65	1.16	6.8
J1246+0238	0.41	>0.68	>1.25	7.3
J1305+5116	0.40	>0.78	>1.56	8.5
J1435+3131	0.52	0.71	1.11	7.5
J1443+4725	0.51	0.86	1.58	7.8
J1505+0326	0.70	>0.86	>1.19	6.6
J1548+3511	0.52	0.80	1.38	7.9
J1633+4718	0.43	0.68	1.19	6.3
J1634+4809	0.44	0.74	1.35	7.4
J1644+2619	0.51	0.74	1.22	6.9
J1722+5654	0.43	0.69	1.24	7.4

^aCalculated using the rest frame luminosities at 5 GHz, 2500 Å, and 2 keV. The lower limits are computed for X-ray non-detections from the X-ray flux limits set by the RASS.

^bVirial black hole masses estimated from the width and luminosity of the broad H β lines (see text for details).